



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/335, C07D 305/14	A1	(11) International Publication Number: WO 97/09979 (43) International Publication Date: 20 March 1997 (20.03.97)
(21) International Application Number: PCT/US96/14613 (22) International Filing Date: 13 September 1996 (13.09.96) (30) Priority Data: 60/003,687 13 September 1995 (13.09.95) US (71) Applicant (for all designated States except US): FLORIDA STATE UNIVERSITY [US/US]; Sliger Building, Room 100, 2035 East Paul Dirac Drive, Tallahassee, FL 32310 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): HOLTON, Robert, A. [US/US]; Dittmer Laboratory of Chemistry, Chemistry Dept., Room 715, Tallahassee, FL 32306 (US). NADIZADEH, Hossain [US/US]; 415 Chapel Drive #D308, Tallahassee, FL 32304 (US). HOFER, Kurt, G. [US/US]; 235 Westridge Drive, Tallahassee, FL 32304 (US). YANG, Li-Xi [US/US]; Apartment 4B, 1839 Miccosukee Road, Tallahassee, FL 32304 (US). (74) Agents: HEJLEK, Edward, J. et al.; Senniger, Powers, Leavitt and Roedel, 16th floor, One Metropolitan Square, St. Louis, MO 63102 (US).		(81) Designated States: AU, CA, CN, CZ, FI, HU, JP, KR, NO, NZ, PL, RU, US, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i> <i>With amended claims.</i>
(54) Title: RADIOSENSITIZING TAXANES AND THEIR PHARMACEUTICAL PREPARATIONS (57) Abstract Taxanes containing electron-affinic radiosensitizing functional groups, their pharmaceutical preparations, and methods of making and using this new class of highly potent radiosensitizers of tumor cells.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgystan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

**RADIOSENSITIZING TAXANES AND
THEIR PHARMACEUTICAL PREPARATIONS**

Background of the Invention

5 This invention relates to novel radiosensitizing compounds, and in particular, to substituted taxanes containing radiosensitizing moieties, their pharmaceutical preparations, and methods of using this new class of highly potent radiosensitizers of tumor cells.

10 In the United States alone, over a half million patients undergo radiation therapy each year as a part of their battle against cancer. To date, however, radiation therapy has produced only limited success as a cancer treatment. Understandably, therefore, a major effort has
15 been underway for a number of years to develop means to improve the efficacy of such radiotherapy techniques.

It is widely believed that the presence of radioresistant, hypoxic (poorly oxygenated) cells in tumors constitutes a significant factor in causing local
20 failure in conventional cancer radiotherapy. For example, it was reported by Gatenby et al., Int. J. Radiat. Oncol. Biol. Phys. 14: 831-833 (1988), that for head and neck tumors, the hypoxic cell volume is inversely correlated with tumor radiosensitivity. Other
25 reports confirm this conclusion for a variety of types of tumors and suggest that the presence of a concentration of as little as 2-3% hypoxic cells in a tumor may double the radiation dose required for tumor control.

Various solutions have been proposed to
30 overcome the problem of hypoxia, including carrying out radiation treatments in high pressure oxygen chambers and the substitution of "fast neutron" or π meson radiation in place of X-rays. However, these techniques are not wholly satisfactory for a number of reasons, including

the great expense and difficulty frequently associated with such procedures.

One promising field of investigation for dealing with radioresistant hypoxic tumor cells has been the use of "radiosensitizing" compounds which selectively increase the sensitivity of hypoxic cells to radiation. This specificity to hypoxic cells is also valuable because a significant percentage of solid tumors are characterized by such cells while most normal tissue is not. Thus, treatment with such compounds serves to enhance the impact of radiation on tumor cells while having little effect on the impact of radiation on healthy cell tissue. A number of heterocyclic, electron-affinic compounds, and in particular, those with oxidized nitrogen moieties, have been successfully used for the purpose of radiosensitizing hypoxic tumor cells. Specifically, the discovery that the nitroimidazoles, metronidazole (metro) and misonidazole (miso), sensitize hypoxic cells to radiation provided initial optimism for a breakthrough solution to the problem of tumor hypoxia. Unfortunately, however, both agents have proven to be highly toxic at therapeutic levels.

The possibility of using chemotherapeutic agents to selectively enhance radiation response in tumors has also been proposed. In addition to its use as a chemotherapeutic agent, taxol, for example, has been investigated in vitro and in vivo as a potential radiosensitizing drug. See, Tishler et al., Radiation Oncology Biol. Phys., 22:613-617 (1992); Tishler et al., Cancer Research, 52:3495-3497 (1992); Steren, et al., Gynecologic Oncology, 48:252-258 (1993); Steren, et al., Gynecologic Oncology, 50:89-93 (1993); Choy et al., Cancer, 71:3774-3778 (1993); Milas et al., Cancer Research, 54:3506-3510 (1994); and Joschko et al., Proceedings of the American Association for Cancer Research, 35:647 (1994). Although the reported data


suggests that taxol is an effective radiosensitizer, recent data from our laboratory leads us to question whether the reported data has been misinterpreted. In any event, a need continues to exist for compounds which
5 possess antitumor activity and which are more potent radiosensitizers and thus, can be administered at lower doses to reduce toxic side effects.

Summary of the Invention

Among the several objects of the invention, therefore, may be noted the provision of a novel class of
10 compounds and pharmaceutical preparations containing them which possess antitumor activity and which are potent radiosensitizing agents for cancer radiation therapy. Also among the objects of the invention are methods for
15 the use of such compounds and pharmaceutical preparations in warm-blooded animals in radiation therapy.

Briefly, therefore, the present invention is directed to taxanes comprising one or more electron-affinic moieties. Such compounds provide greatly
20 enhanced radiosensitization of tumors and reduced toxic side effects to normal body tissues at a given dosage as compared to conventional radiosensitization agents. The electron-affinic moiety may be attached directly, or indirectly through a linker to one of the ring atoms of
25 the taxane or to one of the C13 side chain atoms. For example, the electron-affinic moiety may be attached to the C2, C4, C7, C9, C10, C14, C3' or C5' carbon of a taxane corresponding to the structure:

4



wherein

M comprises ammonium or is a metal;

R₁ is hydrogen or hydroxy;

5 R₂ is -OT₂, -OCOZ₂, -OCOOZ₂, RSG₁ or RSG₂;

R₄ is -OT₄, -OCOZ₄, -OCOOZ₄, RSG₁ or RSG₂;


R₇ is hydrogen, halogen, -OT₇, -OCOZ₇, -OCOOZ₇,
RSG₁ or RSG₂;

10 R₉ is hydrogen, keto, -OT₉, -OCOZ₉, -OCOOZ₉, RSG₁
or RSG₂;

R₁₀ is hydrogen, keto, -OT₁₀, -OCOZ₁₀, -OCOOZ₁₀,
RSG₁ or RSG₂;

R₇, R₉, and R₁₀ independently have the alpha or
beta stereochemical configuration;

15 R₁₃ is hydroxy, protected hydroxy, keto, MO- or



;

R₁₄ is hydrogen, hydroxy, protected hydroxy,
RSG₁ or RSG₂;

20 T₂, T₄, T₇, T₉ and T₁₀ are independently hydrogen
or hydroxy protecting group;

X₁ is -OX₆;

X₂ is hydrogen, hydrocarbon, heterosubstituted
hydrocarbon, heteroaryl or heterosubstituted heteroaryl;

X_3 and X_4 are independently hydrogen, hydrocarbon, heterosubstituted hydrocarbon, heteroaryl, heterosubstituted heteroaryl or RSG_1 ;

X_5 is $-X_{10}$, $-OX_{10}$, $-SX_{10}$, or $-NX_8X_{10}$;

5 X_6 is hydrogen, hydrocarbon, heterosubstituted hydrocarbon, heteroaryl, heterosubstituted heteroaryl or hydroxy protecting group or a functional group which increases the water solubility of the taxane derivative;

10 X_8 is hydrogen, hydrocarbon, heterosubstituted hydrocarbon, RSG_1 or RSG_2 ;

X_{10} is hydrocarbon, heterosubstituted hydrocarbon, heteroaryl, heterosubstituted heteroaryl, RSG_1 or RSG_2 ;

15 Z_2 , Z_4 , Z_7 , Z_9 and Z_{10} are independently hydrocarbon, heterosubstituted hydrocarbon, heteroaryl or heterosubstituted heteroaryl;

RSG_1 is an electron-affinic moiety;

RSG_2 is $-L-(RSG_1)_n$;

20 L is a linker comprising a chain of 1 to 30 atoms in the chain, the atoms being selected from the group consisting of C, O, N, S, Si, and P; and

n is an integer greater than or equal to 1.

The invention is also directed to pharmaceutical compositions for radiosensitizing tumor cells which contain a radiosensitizing amount of the above described taxanes or a pharmaceutically acceptable salt thereof in admixture with a pharmaceutically acceptable carrier.

25 The present invention is further directed to a process for radiosensitizing tumor cells. The process comprises administering a radiosensitizing amount of the pharmaceutical composition described above to the tumor cells. Related thereto, a method is also provided for killing tumor cells in a warm-blooded animal which includes the steps of administering to the warm-blooded animal a pharmaceutical composition as described above in

an amount effective to radiosensitize the tumor cells, followed by, after a time interval sufficient to enhance radiosensitization of the tumor cells, irradiating the tumor cells with a dose of radiation effective to kill the tumor cells.

Other objects and features will be in part apparent and in part pointed out hereinafter.

Brief Description of the Drawings

Fig. 1 is a graph depicting in vitro chemotherapeutic activity of taxoltere metro, taxol and taxoltere pnip on CHO cells for the studies set forth in Example 4.1.

Fig. 2 is a graph depicting in vitro chemotherapeutic activity of taxoltere, taxol and taxoltere pnip on HCT 116 cells for the studies set forth in Example 4.1.

Fig. 3 is a graph depicting in vitro chemotherapeutic radiosensitization of taxoltere metro and taxoltere pnip on CHO cells for the studies set forth in Example 4.2.

Fig. 4 is a graph depicting in vitro chemotherapeutic radiosensitization of taxoltere metro and taxoltere pnip on HCT 116 cells for the studies set forth in Example 4.2.

Fig. 5 is a graph depicting in vivo dose-response curves for taxoltere metro and taxol for the studies set forth in Example 4.4.

Fig. 6 is a graph depicting in vivo chemotherapeutic activity of taxoltere metro and taxoltere pnip at 40% of LD₅₀ for the studies set forth in Example 4.4.

Fig. 7 is a graph depicting the chemotherapeutic effect of low-dose multi-treatment (Q7D x 4) with taxol and its analogs on MTG-B mammary tumors for the studies set forth in Example 4.4.

Fig. 8 is a graph depicting the chemotherapeutic effects of low-dose multi-treatment (Q7D x 4) with taxol and its analogs on MTG-B mammary tumors for the studies set forth in Example 4.4.

5 Figs. 9 and 10 are graphs depicting the chemotherapeutic effects of low-dose multi-treatments (Q11D x 4) with taxol and its analogs on MTG-B mammary tumors for the studies set forth in Example 4.4.

10 Fig. 11 is a graph depicting in vivo chemotherapeutic radiosensitization of taxoltere metro, taxoltere pnip and taxol on MTG-B mammary tumors (i.p., single dose) for the studies set forth in Example 4.5.

15 Fig. 12 is a graph depicting the effects of ip taxol, taxoltere metro and taxoltere pnip +/-RT on MTG-B mammary tumors for the studies set forth in Example 4.5.

Figs. 13 and 14 are graphs depicting the effects of taxoltere pnip on MTG-B mammary tumors (i.v., single dose, 24% LD50) for the studies set forth in Example 4.5.

20 Fig. 15 is a graph depicting the cure rate for taxoltere pnip (i.p., single dose) for the studies set forth in Example 4.6.

25 Figs. 16 and 17 are graphs depicting the cure rate for taxoltere pnip (i.v., single dose) for the studies set forth in Example 4.6.

Fig. 18 is a graph depicting the cure rate for taxoltere pnip +/- RT on MTG-B mammary tumors in vivo as a function of drug dose for the studies set forth in Example 4.6.

30 Detailed Description of the Preferred Embodiments

Surprisingly, it has been discovered that taxanes containing electron-affinic substituents exhibit significantly greater potency than taxol as a radiosensitizing agent. As a result, such increased
35 potency permits the administration of much lower dosages


of these compounds for the same or even greater radiosensitization of tumor cells, allowing for a concomitant reduction in toxic side effects on healthy tissue for any particular dosage level required to effectively radiosensitize the tumor cells.

The radiosensitizing groups or moieties described herein (RSG₁ and RSG₂) impart electron affinity to the compounds with which they are associated. This novel class of potent radiosensitizers comprise taxanes containing at least one, and optionally two or more electron-affinic moieties. In general, the radiosensitizing moieties contain electron-affinic groups which fall into one of four groups: (i) carbocyclic or heterocyclic aromatic moieties which possess one or more carbonyl, trifluoromethyl, halogen, nitro, sulfonyl, sulfinyl, phosphoryl, oxide or cyano groups, (ii) heterocyclic aromatic moieties containing two or more heteroatoms, (iii) metal complexes, and (iv) organo-metallic groups in which the metal is covalently bonded to carbon.

The carbocyclic or heterocyclic aromatic electron-affinic moieties contain one to three rings with a total of 5 to 15 ring atoms which are selected from the group consisting of C, N, S, O and P. Preferably, the carbocyclic or heterocyclic aromatic electron-affinic moieties contain one to two rings with one ring being presently most preferred. Representative carbocyclic aromatic electron-affinic moieties include phenyl and naphthyl groups containing one or more nitro, halogen, carbonyl or sulfonyl substituents, with nitro-substituted phenyl being a preferred carbocyclic aromatic electron-affinic moiety. Representative heterocyclic aromatic electron-affinic moieties include imidazoles, triazoles, pyridines, benzamides, nicotinamides, benzotriazine oxides, furans, thiophenes, oxazoles and thiozoles possessing one or more carbonyl, trifluoromethyl,

halogen, nitro, sulfonyl, sulfinyl, phosphoryl, oxide or cyano groups, and preferably at least one nitro group.

Nitroimidazole and nitrotriazole heterocyclic aromatic electron-affinic moieties which may be
 5 incorporated into the radiosensitizing agents of the present invention include 2-nitroimidazol-1-yl and 3-nitro-1,2,4-triazol-1-yl and other nitroimidazoles and nitrotriazoles which correspond to the following structures:



wherein E₁ is alkyl or fluoroalkyl. The preparation and use of radiosensitizing agents incorporating these and other nitroimidazoles and nitrotriazoles is described in Suzuki et al., U.S. Patent Nos. 4,945,102 and 5,064,849;
 15 Kagiza et al., U.S. Patent Nos. 4,927,941, 4,977,273 and 5,304,654; Suto, U.S. Patent No. 4,954,515 and 5,036,096; Suto et al., U.S. Patent No. 4,797,397; Papadopolou-Rosenzweig et al., U.S. Patent No. 5,294,715; Beylin et al., U.S. Patent No. 5,342,959.


20 Benzamide and nicotinamide heterocyclic aromatic electron-affinic moieties which may be incorporated into the radiosensitizing agents of the present invention include

- 25 5-hydroxynicotinamide;
- 5-nitronicotinamide;
- 5-(2,3-dihydroxypropoxy)nicotinamide;
- 5-aminonicotinamide;
- 5-(2-methoxyethylamino)nicotinamide;
- 5-acetamidonicotinamide;
- 30 3-hydroxy thiobenzamide;

10

3-[(2-hydroxyethoxy)acetamido]benzamide;
 3-(2,3 dihydroxy-n-propoxy)-4-methoxybenzamide;
 3-(2,3 dihydroxy-n-propoxy)-4-methylbenzamide;
 4-(2,3 dihydroxy-n-propoxy)-3-methoxybenzamide;

- 5 and other benzamides and nicotinamides which correspond to the following structures:




- wherein X^1 is O or S; Y_1 is H, lower alkyl, lower alkoxy, acetoxyl, or acetamido; Y_2 is -OR, -SR, -NHR, -NO₂,
 10 -O(CO)R, -NH(CO)R, -O(SO)R, or -O(POR)R; Y_3 is H, Z_1 , -OR, -SR, -NHR, -O(CO)R, -NH(CO)R, -O(SO)R, or -O(POR)R; and R is hydrogen or hydrocarbon which may be optionally substituted and interrupted by an ether (-O-) linkage.
 The preparation and use of radiosensitizing agents
 15 incorporating these and other benzamides and nicotinamides is described in Lee et al., U.S. Patent Nos. 5,032,617, 5,041,653 and 5,175,287.

- 20 Benzotriazine oxide heterocyclic aromatic electron-affinic moieties which may be incorporated into the radiosensitizing agents of the present invention include


- 3-hydroxy-1,2,4-benzotriazine-1,4-dioxide;
 3-amino-7-trifluoro-1,2,4-benzotriazine-1-oxide;
 25 3-amino-7-decyl-1,2,4-benzotriazine-1-oxide;
 3-amino-7-carbamyl-1,2,4-benzotriazine-1-oxide;
 7-acetyl-3-amino-1,2,4-benzotriazine-1-oxide;
 7-chloro-3-hydroxy-1,2,4-benzotriazine-1,4-dioxide;

7-nitro-3-amino-1,2,4-benzotriazine-1,4-dioxide; and other benzotriazine oxides corresponding to the structure:



- 5 wherein Y_4 is H, substituted or unsubstituted lower hydrocarbon, or alkanoyl; m is 0 or 1; and Y_5 and Y_6 are independently hydrogen, nitro, halogen, morpholino, pyrrolidino, piperidino, substituted or unsubstituted hydrocarbon, $-NH_2$, $-NHR'$, $-NR'R'O(CO)R'$, $-NH(CO)R'$,
 10 $-O(SO)R'$, or $-O(POR')R'$ in which R' is substituted or unsubstituted hydrocarbon. The preparation and use of radiosensitizing agents incorporating these and other benzotriazine oxides is described in Lee et al. U.S. Patent No. 5,175,287.
- 15 The metal complex electron-affinic moieties preferably comprise Pt^{2+} , Co^{3+} , Fe^{2+} , Fe^{3+} , Pd^{2+} , Cu^{2+} , Ti^{4+} , or Zr^{4+} as the metal and generally fall into two subgroups: (a) metal complexes of the carbocyclic and heterocyclic aromatic electron-affinic moieties discussed
 20 above, and (b) metal complexes of bidentate ligands comprising nitrogen, carbon or sulfur. In general, metal complexes of bidentate ligands correspond to the formula $-BM^LX_k$ wherein B is a bidentate ligand containing nitrogen, carbon or sulfur, M^L is a metal, X is an anionic
 25 ligand such as Cl^- or OAc^- , and k is 1-4. Exemplary bidentate ligands include:

12



Electron-affinic metal complexes which may be incorporated into the radiosensitizing agents of the present invention include compounds of the formula:


5 $[PtX^M_j(NR_2''H)Q]$ or $[PtX^M_j(NR_2''H)_2Q]^+Y^-$

wherein n is 1 or 2, and wherein when n is 2, X^M is a monovalent biologically acceptable anion, and when j is 1, X^M is a divalent biologically acceptable anion; each R'' is independently H or alkyl, or both R'' s together are a

10 piperidino or morpholino residue; Q is a radiosensitizing ligand selected from a mononitro-substituted imidazole, a mononitro-substituted pyrazole, a mononitro-substituted thiazole and a mononitro-substituted isothiazole; and Y^- is a physiologically acceptable anion. These

15 heterocycles may optionally be substituted by an alkyl, amino substituted alkyl, hydroxy, alkoxy or amino group. In addition, if the heterocycle is pyrazole or imidazole, a ring nitrogen may be substituted by alkyl or alkoxy or hydroxy substituted alkyl and wherein one or two

20 methylenes of the alkyl may be replaced by oxygen. In a preferred embodiment, Q is one of the following:




wherein R^1 is alkyl optionally containing an amino substituent, $-OR^3$, or $-N(R^3)_2$, wherein R^3 is H or lower

25 alkyl; R^2 is alkyl or 1-8 carbons substituted by one or more $-OR^3$ and wherein one or two methylenes may be

replaced by oxygen and each m is independently 0 or 1. The preparation and use of radiosensitizing agents incorporating these metal complexes is described in Skov et al. U.S. Patent Nos. 4,921,963 and 5,026,694.


5 Other electron-affinic metal complexes which may be incorporated into the radiosensitizing agents of the present invention may be made by reacting an organic or inorganic platinum compound such as an alkali metal tetrahaloplatinate or cis-bis(acetonitrile)dichloro-
10 platinum (II) with rhodamine 123 or other (+)-charged rhodamine or the like, for example, a cyanine dye such as 3,3'-diethylthiadicarbocyanine iodide or other (+)-charged cyanine dyes as described in U.S. Patent No. 5,196,413.

15 Other electron-affinic metal complexes which may be incorporated into the radiosensitizing agents of the present invention include include Cu(II) compounds selected from compounds having the formula:



20 wherein A^c represents a bidentate heteroaromatic ligand containing neutral nitrogen donor atoms; B^c represents a bidentate ligand containing neutral or negatively charged oxygen donor atoms; X^c and Y^c are the same or different neutral or negatively charged monodentate ligands; and Z¹
25 and Z² represent the charge on the complex. The preparation and use of radiosensitizing agents incorporating these metal complexes is described in Abrams et al. U.S. Patent No. 5,100,885.

30 Other electron-affinic metal complexes which may be incorporated into the radiosensitizing agents include Co(III) or Fe(III) compounds a formula corresponding to one of the following formulas:



35 wherein n has a value of 3 or 4; N is an uncharged nitrogen donor atom that is contained within a ligand; X^F represents an anionic ligand; and y represents the charge

on the complex; A^F represents a bidentate or tetradentate negative ligand containing N or O donor atoms; D^1 and D^2 represent the same or different monodentate ligands; q represents a positive or negative charge on the complex;
5 Z^F represents a chelating mononegative negative ligand; T^1 and T^2 , which may be the same or different, represent mono-negative tridentate ligands. The preparation and use of radiosensitizing agents incorporating these metal complexes is described in U.S. Patent No. 4,727,068.

10 The organometallic electron-affinic moieties are aliphatic or aromatic mercury radicals. The preparation and use of radiosensitizing agents incorporating mercury containing entities is described in Shenoy et al., Cancer Investigation, 10(6):533-551 (1992)
15 and Bruce et al., Radiation Res., 24:473-481 (1965).







The electron-affinic moieties may be directly attached to one of the carbons of the A, B, or C rings of the taxane or indirectly attached via a linker. The linker comprises a chain of 0 to 30 atoms in the chain,
20 with approximately 10 or less being preferred. The chain atoms are selected from the group consisting of C, O, N, S, Si, and P and are preferably C, N or O. The linker may be linear or cyclic, branched or unbranched, and may contain as substituents, one or more P, C, O, N, S, H, Si
25 or halogen-containing substituents. Exemplary linker substituents include silyls, ethers, thioethers, esters, thioesters, amides, thioamides, amines, alcohol, alkyl, aryl, carbonyl, sulfonyl, phosphoryl, and halogen substituents.

30 Preferably, the linker comprises a hydrocarbon segment consisting of 1 to 6 carbon atoms. It may additionally comprise a carbonyl, ester, thioester, amide, carbonate, thiocarbonate, carbamate, or ether segment. If a non-hydrocarbon segment is included; the
35 non-hydrocarbon segment preferably comprises one or more

15





ether, carbonate or carbonyl moieties as the non-hydrocarbon segment.

For purposes of illustration, a series of radicals comprising linkers and electronic-affinic moieties falling within the scope of the present invention is set forth as follows:



10

16



5


wherein h is 1-3, R⁴ is H, hydrocarbon or substituted hydrocarbon, and R⁵ is hydrocarbon or substituted hydrocarbon. In other embodiments, the carbonyl or ester linkage of the above structures may be replaced by

10 thioester or amide linkages. In addition, many of these


radicals may serve as ligands for the previously identified metal species.

The radiosensitizing compounds of the present invention are prepared by linking the electron-affinic moiety to the C2, C4, C7, C9, C10, C14, C3', or C5' carbons of a taxane. The starting material may be 10-deacetyl baccatin III, baccatin III, or another naturally occurring taxane such as 14-hydroxy-10-deacetyl baccatin III. Alternatively, the taxane may be synthesized from commodity chemicals as set forth in PCT Patent Application No. WO 95/03265.

Taxanes having C13 side chains which incorporate electron-affinic moieties at C3' and/or C5' may be prepared through the use of β -lactams having the desired substituents and reacting the β -lactam and a C13 metal or ammonium alkoxide of a suitably substituted taxane as more fully described in U.S. Patent 5,430,160. The β -lactams have the following structural formula:



wherein X_1 is protected hydroxy, and $X_2 - X_5$ are as previously defined. Preferably, the alkoxide has the tetracyclic taxane nucleus and corresponds to the structural formula:




wherein M is a metal or tetraalkylammonium and R₁, R₂, R₄, R₇, R₉, R₁₀, and R₁₄ are as previously defined.

5 The electron-affinic moieties can be attached to the C2, C4, C7, C9, C10, and C14 positions of a taxane by a variety of methods. For purposes of illustration, the attachment method will first be described with respect to the C7 position. As will be described elsewhere herein, however, these same methods can be used for the other positions.





10 Metronidazole, a well known radiosensitizer, and other electron-affinic moieties can be attached via a carbonate linkage by treating baccatin III 2 with carbonyl diimidazole to produce the 7-carboxylimidazolide 2a, and reacting the product in situ with metronidazole
15 at higher temperature to provide the 7-carboxylmetronidazolide 3. A C13 side chain can then be attached to 7-carboxylmetronidazolide 3 by treating it with lithium hexamethyldisilazide and β -lactam 4, and, after treatment with HF, to yield a taxane which we have
20 named taxoltere metro 5. This reaction sequence is summarized in the following reaction scheme:

19




This method can be used for the preparation of a series of similar radiosensitizing taxanes having different radiosensitizing groups. The intermediate 7-carboxylimidazolidine **2a** reacts smoothly with alcohols to provide the desired radiosensitizing taxanes in which the radiosensitizing group is attached via a carbonate linkage, as in taxoltere metro. For example, alcohols **6**

through 9 will react with the carbonylimidazolidine substituent of 2a to yield four other taxanes having radiosensitizing groups linked to C7.




- 5 Attachment of the radiosensitizing group is then followed by attachment of a side chain at C13 in the same manner as it was accomplished for taxoltere metro.


The attachment of a metal atom or metal complex tethered at the C7 position of the taxane core can be accomplished by reacting an allylchloroformate with a taxane having an available C7 hydroxy group and a protected 2' hydroxy group to produce derivative 10 wherein X₃ and X₅ are as previously defined and P is a hydroxy protecting group. Hydroboration of the allyl carbonate substituent followed by treatment of the borane with mercuric acetate and sodium chloride and deprotection of the C2' hydroxy group gives the mercury derivative 11, a good radiosensitizer.




21





The ester analog of 11 can be prepared by the direct acylation of the C7 hydroxyl group of a 2' hydroxy protected taxane with an acid chloride to produce ester 12. Hydroboration of the allyl ester followed by treatment of the borane with mercuric acetate and sodium chloride gives the ester analog of mercury derivative 11.




Similar chemistry can be used to attach a bidentate ligand to the C7 position, and the metal complex (e.g., platinum) of the bidentate ligand can then be prepared by introducing an appropriate metallic reactant (e.g., PtCl_2 or $\text{PtCl}_2(\text{SMe}_2)_2$). The bidentate ligand can also incorporate an electron-affinic ligand, and the Pt (II) complex of 13 can be made in this way.




The length and nature of the linker between the taxane and the electron-affinic group may be altered. It is possible to prepare analogs in which the electron-affinic group is both closer and further away from the taxane than it is in taxoltere metro. An analog with a longer linker can easily be synthesized by incorporating a dicarboxylic acid diester instead of the carbonate between the taxane and metronidazole, e.g., 14. The *p*-nitrobenzyl ether 15 and the corresponding *p*-nitrobenzoate are radiosensitizing taxanes in which the electron-affinic group is very close to the taxane. Alternatively, hydroboration of 10 gives an alcohol, the mesylate of which reacts with, for example, 2-nitroimidazole to provide 16, and the epoxide derived by peracid treatment of 10 reacts with, for example, 2-nitroimidazole to provide 17. Ester analogs of 16 and 17 can be similarly prepared starting from 12.




23



It is also possible to prepare radiosensitizing taxanes having multiple radiosensitizing groups attached to a single linker. For example, reaction of 2-nitro
 5 imidazole with glycidyl chloride at somewhat elevated temperature provides alcohol 18, which then reacts with 2a and with β -lactam 4 to give radiosensitizing taxane 19.



10



Electron-affinic moieties can be attached to the C10 position of a taxane possessing a C10 hydroxy group, such as 10-DAB, by the methods discussed for

5 attaching the electron-affinic moieties to the C7 position. Taxanes having other C10 substituents described herein may be prepared as more fully described in PCT Patent Application WO 94/15599 and other literature references. For example, taxanes having a C10

10 keto substituent can be prepared by oxidation of 10-desacetyl taxanes. Taxanes which are dihydro substituted at C10 can be prepared by reacting a C10 hydroxy or acyloxy substituted taxane with samarium diiodide. Taxanes having acyloxy substituents other than acetate

15 can be prepared by reacting the C10 hydroxy substituent of 10-deacetyl baccatin III with any standard acylating agent such as anhydrides, acid chlorides, acyl imidazoles or other activated carboxyl derivatives. Taxanes having a C10 carbonate substituent can be prepared by using an

20 analogous chloroformate instead of the acid chloride.

Electron-affinic moieties can be attached to the C9 position of a taxane possessing a C9 hydroxy group by the methods discussed for attaching the electron-affinic moieties to the C7 position. As more fully


25 described in PCT Patent Application WO 94/20088, the C9 the C9 keto substituent of taxol, 10-DAB, baccatin III or can be selectively reduced to yield the corresponding C9

β -hydroxy derivative with a borohydride, preferably tetrabutylammonium borohydride (Bu_4NBH_4) or triacetoxy-borohydride. The C9 β -hydroxy derivative can then be protected at C7 with a hydroxy protecting group and the C9 hydroxy group can be acylated following the methods described herein for acylation of the C7 hydroxy group.

Alternatively, reaction of 7-protected-9 β -hydroxy derivative with KH causes the acetate group (or other acyloxy group) to migrate from C10 to C9 and the hydroxy group to migrate from C9 to C10, thereby yielding a 10-desacetyl derivative, which can be acylated as described elsewhere herein.

As more fully described in PCT Patent Application WO 94/17050, C7 dihydro and other C7 substituted taxanes can be prepared by tin hydride reduction of the C7 xanthate. C7 fluoro-substituted taxanes can be prepared by treatment of C13-triethylsilyl-protected baccatin III with 2-chloro-1,1,2-trifluorotriethylamine at room temperature in THF solution. Other baccatin derivatives with a free C7 hydroxyl group behave similarly. Alternatively, 7-chloro baccatin III can be prepared by treatment of baccatin III with methanesulfonyl chloride and triethylamine in methylene chloride solution containing an excess of triethylamine hydrochloride. Taxanes having C7 acyloxy substituents can be prepared as set forth in the following reaction scheme. 7,13-protected 10-oxo-derivative is converted to its corresponding C13 alkoxide by selectively removing the C13 protecting group and replacing it with a metal such as lithium. The alkoxide is then reacted with a β -lactam or other side chain precursor. Subsequent hydrolysis of the C7 protecting groups causes a migration of the C7 hydroxy substituent to C10, migration of the C10 oxo substituent to C9, and migration of the C9 acyloxy substituent to C7.

26




Taxanes having alternative C2 and/or C4 esters or carbonates which optionally may contain an electron-affinic moiety as described elsewhere herein can be prepared using baccatin III and 10-DAB as starting materials. The C2 and/or C4 esters of baccatin III and 10-DAB can be selectively reduced to the corresponding alcohol(s) using reducing agents such as LAH or Red-Al, and new esters can thereafter be substituted using standard acylating agents such as anhydrides and acid chlorides in combination with an amine such as pyridine, triethylamine, DMAP, or diisopropyl ethyl amine.

Alternatively, the C2 and/or C4 alcohols may be converted to new C2 and/or C4 esters through formation of the corresponding alkoxide by treatment of the alcohol with a suitable base such as LDA followed by an acylating agent such as an acid chloride. The corresponding carbonates can be prepared by substituting a chloroformate for the analogous acid chloride.


Baccatin III and 10-DAB analogs having different substituents at C2 and/or C4 can be prepared as set forth in Reaction Schemes C₂-1 to C₂-5. To simplify the description, 10-DAB is used as the starting material and only the ester products are shown. It should be understood, however, that other starting materials and reactants may be substituted to yield the other C2 and C4 substituted compounds disclosed herein.

In the Reaction Scheme C₂-1, protected 10-DAB 20 is converted to the triol 21 with lithium aluminum hydride. Triol 21 is then converted to the corresponding C4 ester using Cl₂CO in pyridine followed by a nucleophilic agent (e.g., Grignard reagents or alkyllithium reagents) wherein Z₂ is as defined elsewhere herein.

28


Reaction Scheme C₂-1

Deprotonation of triol 21 with LDA followed by introduction of an acid chloride selectively gives the C4 ester. For example, when acetyl chloride was used, triol 21 was converted to 1,2 diol 24 as set forth in Reaction Scheme C₂-2 wherein Z₄ is as defined elsewhere herein.


Reaction Scheme C₂-2

Triol **21** can also readily be converted to the 1,2 carbonate **22**. Acetylation of carbonate **22** under vigorous standard conditions provides carbonate **25** as described in Reaction Scheme C₂-3; addition of alkylolithiums or Grignard reagents to carbonate **22** provides the C2 ester **24** having a free hydroxyl group at C4 as set forth in Reaction Scheme C₂-1.


Reaction Scheme C₂-3




As set forth in Reaction Scheme C₂-4, other C4 substituents can be provided by reacting carbonate **22** with an acid chloride and a tertiary amine to yield carbonate **26** which is then reacted with alkylolithiums or Grignard reagents to provide 10-DAB derivatives **27** having new substituents at C2 wherein Z₂ and Z₄ are as defined elsewhere herein.

Reaction Scheme C₂-4

Alternatively, baccatin III may be used as a starting material and reacted as shown in Reaction Scheme C₂-5. After being protected at C7 and C13, baccatin III is reduced with LAH to produce 1,2,4,10 tetraol 29. Tetraol 29 is converted to carbonate 30 using Cl_2CO and pyridine, and carbonate 30 is acylated at C10 with an acid chloride and pyridine to produce carbonate 31 (as shown) or with acetic anhydride and pyridine (not shown). Acetylation of carbonate 31 under vigorous standard conditions provides carbonate 32 which is then reacted with alkyl lithiums to provide the baccatin III derivatives 33 having new substituents at C2 and C10 wherein Z_2 and Z_{10} are as defined elsewhere herein.

Reaction Scheme C₂-5

32



Taxanes having radiosensitizing groups at the C14 position, i.e., the point of attachment of R₁₄ as depicted elsewhere herein, may be prepared using the same or similar methods as those described elsewhere herein with respect to attaching radiosensitizing groups to the C7 position of the taxane. The starting material for these compounds may be, for example, a hydroxylated taxane (14-hydroxy-10-deacetylbaccatin III) which has been discovered in an extract of yew needles (C&EN, p 36-37, April 12, 1993). Derivatives of this hydroxylated taxane having the various C2, C4, C7, C9, C10, C3' and C5' functional groups described above may also be prepared by using this hydroxylated taxane. In addition, the C14 hydroxy group together with the C1 hydroxy group of 10-DAB can be converted to a 1,2-carbonate as described in C&EN or it may be converted to a variety of

esters or other functional groups as otherwise described herein in connection with the C2, C4, C9 and C10 substituents.

The taxane radiosensitizers of the present invention can be combined with various excipient vehicles and/or adjuvants well known in this art which serve as pharmaceutically acceptable carriers to permit drug administration in the form of, e.g., injections, suspensions, emulsions, tablets, capsules, and ointments. These pharmaceutical compositions, containing a radiosensitizing amount of the described substituted diamine compounds, may be administered by any acceptable means which results in the radiosensitization of tumor cells. For warm-blooded animals, and in particular, for humans undergoing radiotherapy treatment, administration can be oral, parenteral, subcutaneous, intravenous, intramuscular and/or intraperitoneal. To destroy tumor cells, the pharmaceutical composition containing the radiosensitizing diamines are administered in an amount effective to radiosensitize the tumor cells (in the range of 1 to 100 mg/kg for humans). The specific dosage administered will be dependent upon such factors as the general health and physical condition of the patient as well as his age and weight, the stage of the patient's disease condition, and the existence of any concurrent treatments.

After administration of the radiosensitizing composition to the tumor cells and the passage of a time interval sufficient to enhance radiosensitization of the tumor cells, the tumor cells are irradiated with a dose of radiation effective to destroy the tumor cells. Generally, the patient will receive a total radiation dosage of about 60 to 76 Gy over seven to eight weeks, each individual radiation dose to be given within approximately 1 to 4 hours after administration of the radiosensitizer. Such sequences of radiosensitization

treatments and irradiation are repeated as needed to abate and, optimally, reduce or eliminate, the spread of the malignancy.

Definitions

5 The "hydrocarbon" moieties described herein are organic compounds or radicals consisting exclusively of the elements carbon and hydrogen. These moieties include alkyl, alkenyl, alkynyl, and aryl moieties. These moieties also include alkyl, alkenyl, alkynyl, and aryl
10 moieties substituted with other aliphatic or cyclic hydrocarbon groups, such as alkaryl, alkenaryl and alkynaryl. Preferably, these moieties comprise 1 to 20 carbon atoms.

 The alkyl groups described herein are
15 preferably lower alkyl containing from one to six carbon atoms in the principal chain and up to 20 carbon atoms. They may be straight or branched chain and include methyl, ethyl, propyl, isopropyl, butyl, hexyl and the like. They may be substituted with aliphatic or cyclic
20 hydrocarbon radicals or hetero-substituted with the various substituents defined herein.

 The alkenyl groups described herein are preferably lower alkenyl containing from two to six carbon atoms in the principal chain and up to 20 carbon
25 atoms. They may be straight or branched chain and include ethenyl, propenyl, isopropenyl, butenyl, isobutenyl, hexenyl, and the like. They may be substituted with aliphatic or cyclic hydrocarbon radicals or hetero-substituted with the various substituents
30 defined herein.

 The alkynyl groups described herein are preferably lower alkynyl containing from two to six carbon atoms in the principal chain and up to 20 carbon
35 atoms. They may be straight or branched chain and include ethynyl, propynyl, butynyl, isobutynyl, hexynyl,

and the like. They may be substituted with aliphatic or cyclic hydrocarbon radicals or hetero-substituted with the various substituents defined herein.

The aryl moieties described herein contain from
5 6 to 20 carbon atoms and include phenyl. They may be hydrocarbon or heterosubstituted with the various substituents defined herein. Phenyl is the more preferred aryl.


The heteroaryl moieties described are
10 heterocyclic compounds or radicals which are analogous to aromatic compounds or radicals and which contain a total of 5 to 20 atoms, usually 5 or 6 ring atoms, and at least one atom other than carbon, such as furyl, thienyl, pyridyl and the like. The heteroaryl moieties may be
15 substituted with hydrocarbon, heterosubstituted hydrocarbon or hetero-atom containing substituents with the hetero-atoms being selected from the group consisting of nitrogen, oxygen, silicon, phosphorous, boron, sulfur, and halogens. These substituents include lower alkoxy
20 such as methoxy, ethoxy, butoxy; halogen such as chloro or fluoro; ethers; acetals; ketals; esters; heteroaryl such as furyl or thienyl; alkanoxy; hydroxy; protected hydroxy; acyl; acyloxy; nitro; amino; and amido.

The heterosubstituted hydrocarbon moieties
25 described herein are hydrocarbon moieties which are substituted with at least one atom other than carbon, including moieties in which a carbon chain atom is substituted with a hetero atom such as nitrogen, oxygen, silicon, phosphorous, boron, sulfur, or a halogen atom.
30 These substituents include lower alkoxy such as methoxy, ethoxy, butoxy; halogen such as chloro or fluoro; ethers; acetals; ketals; esters; heteroaryl such as furyl or thienyl; alkanoxy; hydroxy; protected hydroxy; acyl; acyloxy; nitro; amino; and amido.

The acyl moieties described herein contain hydrocarbon, substituted hydrocarbon or heteroaryl moieties.

The alkoxy-carbonyloxy moieties described herein
5 comprise lower hydrocarbon or substituted hydrocarbon moieties.

The term "taxane" as used herein, denotes compounds containing the A, B and C rings (with numbering of the ring positions shown herein):



10

As used herein "Ac" means acetyl; "AIBN" means
azo-(bis)-isobutyronitrile; "Ar" means aryl; "BMDA" means
BrMgNiPr₂; "BOC" means butyloxycarbonyl; "BOM" means
benzyloxymethyl; "10-DAB" means 10-desacetylbaccatin III;
15 "DBU" means diazabicycloundecane; "DMAP" means
p-dimethylamino pyridine; "DDQ" means dicyano-
dichloroquinone; "DMF" means dimethylformamide; "Et"
means ethyl; "FAR" means 2-chloro-1,1,2-trifluoro-
triethylamine; "iPr" means isopropyl; "LAH" means lithium
20 aluminum hydride; "LDA" means lithium diisopropylamide;
"LHMDS" means lithium hexamethyldisilazide; "LTMP" means
lithium tetramethylpiperidide; "mCPBA" means metachloro-
perbenzoic acid; "Me" means methyl; "MOP" means 2-
methoxy-2-propyl; "Ms" means CH₃SO₂-; "Ph" means phenyl;
25 "protected hydroxy" means -OP or -OT wherein P or T is a
hydroxy protecting group; "py" means pyridine; "R" means
lower alkyl unless otherwise defined; "Red-Al" means
sodium bis(2-methoxyethoxy) aluminum hydride; "Swern"
means (COCl)₂, Et₃N; "TASF" means tris(diethylamino)-
30 sulfoniumdifluorotrimethyl-silicate; "TBAF" means
tetrabutylammonium fluoride; "tBu" and "t-Bu" means tert-

butyl; "TBS" means $\text{Me}_2\text{t-BuSi-}$; "TES" means triethylsilyl; "Tf" means $-\text{SO}_2\text{CF}_3$; "TMS" means trimethyl-silyl; "TPAP" means tetrapropylammonium perruthenate; and "Ts" means toluenesulfonyl. "Hydroxy protecting group" includes, but is not limited to, acetals having two to ten carbons, ketals having two to ten carbons, ethers such as methyl, t-butyl, benzyl, p-methoxybenzyl, p-nitrobenzyl, allyl, trityl, methoxymethyl, methoxyethoxymethyl, ethoxyethyl, tetrahydropyranyl, tetrahydrothiopyranyl, and trialkylsilyl ethers such as trimethylsilyl ether, triethylsilyl ether, dimethylarylsilyl ether, triisopropylsilyl ether and t-butyl dimethylsilyl ether; esters such as benzoyl, acetyl, phenylacetyl, formyl, mono-, di-, and trihaloacetyl such as chloroacetyl, dichloroacetyl, trichloroacetyl, trifluoroacetyl; and carbonates including but not limited to alkyl carbonates having from one to six carbon atoms such as methyl, ethyl, n-propyl, isopropyl, n-butyl, t-butyl; isobutyl, and n-pentyl; alkyl carbonates having from one to six carbon atoms and substituted with one or more halogen atoms such as 2,2,2-trichloroethoxymethyl and 2,2,2-trichloroethyl; alkenyl carbonates having from two to six carbon atoms such as vinyl and allyl; cycloalkyl carbonates having from three to six carbon atoms such as cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl; and phenyl or benzyl carbonates optionally substituted on the ring with one or more C_{1-6} alkoxy, or nitro. Other hydroxyl protecting groups may be found in "Protective Groups in Organic Synthesis" by T. W. Greene, John Wiley and Sons, 1981.

To further illustrate and explain the invention, several examples are presented below.

EXAMPLE 1


Preparation of 7-(metronidazoleoxycarbonyl)baccatin III
hn-4-99-4d-2-1

To a solution of baccatin III (100 mg, 0.170
5 mmol) in anhydrous 1,2-dichloroethane (1 mL) under
nitrogen was added 1,1'-carbonyldiimidazole (55 mg, 0.40
mmol) and the reaction mixture was warmed up to 60 °C and
stirred at that temperature for 10 h at which time the
reaction was complete. Metronidazole (292 mg, 1.72 mmol)
10 was added (neat) and the mixture was refluxed under
nitrogen atmosphere. The progress of the reaction was
monitored by NMR. When the reaction was complete
(approximately three days) the mixture was diluted with
ethyl acetate and washed with saturated sodium
15 bicarbonate, brine. The organic layer was separated,
dried and concentrated. The crude mixture was purified
by flash chromatography to give 81 mg (70%) of the 7-
(metronidazole-oxycarbonyl)baccatin III: mp. 220-223
°C; $[\alpha]_{\text{Hg}}^{25} = -44.9$

20 ^1H NMR (CDCl_3 , 500 MHz) δ 8.10 (d, $J=7.1$ Hz, 2H,
benzoate), 7.96 (s, 1H, imidazole), 7.62-7.46 (m, 3H,
benzoate), 6.28 (s, 1H, H10), 5.61 (d, $J=6.8$ Hz, 1H,
H2 β), 5.49 (m, 1H, H7) 4.96 (d, $J=8.2$ Hz, 1H, H5), 4.85
(br s, 1H, H13) 4.62-4.43 (m, 4H, methylenes), 4.31 (d,
25 $J=8.5$ Hz, 1H, H20 α), 4.13 (d, $J=7.5$ Hz, 1H, H20 β), 3.99
(d, $J=6.8$ Hz, 1H, H3), 2.59 (m, 1H, H6 α), 2.50 (s, 3H,
CH3-imidazole), 2.29 (m, 5H, 4Ac, H14's), 2.13 (s, 3H,
10Ac), 2.10 (br s, 3H, Me18) 2.06 (d, $J=5$ Hz, 1H, 13OH),
1.9 (m, 1H, H6 β), 1.77 (s, 3H, Me19), 1.59 (s, 1H, 1OH),
30 1.17 (s, 3H, Me17), 1.08 (s, 3H, Me16).

EXAMPLE 2

Preparation of N-debenzoyl-N-(t-butylcarbamoyl)-7-(metronidazoleoxycarbonyl) taxol. (Taxoltere metro)



5 hn-4-119-2

To a solution of 7-(metronidazoleoxycarbonyl) baccatin III (31 mg, 0.039 mmol) in 0.3 mL of THF at -45 °C was added dropwise 0.048 mL of a 1.00 M solution of lithium bis(trimethylsilyl)amide in THF. After 0.5 h at
 10 -45 °C, a solution of *cis*-1-(t-butoxycarbonyl)-3-triethylsilyloxy-4-phenylazetidin-2-one (75 mg, 0.20 mmol) in 0.3 mL of THF was added dropwise to the mixture. The solution was warmed to 0 °C and kept at that temperature for 1 h before 0.2 mL of a 10% solution of
 15 AcOH in THF was added. The mixture was partitioned between saturated aqueous NaHCO₃ and 60/40 ethyl acetate/hexane. Evaporation of the organic layer gave a residue which was purified by filtration through silica gel to give 45.0 mg of a mixture containing (2'R,3'S)-2'-
 20 triethylsilyl-N-debenzoyl-N-(t-butylcarbamoyl)-7-(metronidazoleoxycarbonyl) taxol and a small amount of the (2'S,3'R) isomer.


To a solution of 45.0 mg of the mixture obtained from the previous reaction in 1.5 mL of
 25 acetonitrile and 0.6 mL of pyridine at 0 °C was added 0.2 mL of 48% aqueous HF. The mixture was stirred at 0 °C for 3 h, then at 25 °C for 13 h, and partitioned between saturated aqueous sodium bicarbonate and ethyl acetate. Evaporation of the ethyl acetate solution gave 40.8 mg of

material which was purified by plug filtration and
 recrystallization from methanol/water to give 32.3 mg
 (79%) of N-debenzoyl-N-(t-butylcarbamoyl)-7-
 (metronidazoleoxy-carbonyl) taxol. m.p. 169-172 °C; $[\alpha]^{25}_{Na}$
 5 -52 °C (0.0035, CHCl₃).

¹H NMR (CDCl₃, 300 MHz) δ 8.10 (d, J=7.1 Hz, 2H,
 benzoate), 7.96 (s, 1H, imidazole), 7.62-7.26 (m, 8H,
 benzoate, 3'-phenyl), 6.25 (s, 1H, H10), 6.18 (dd, J = 8.8,
 8.8 Hz, 1H, H13), 5.65 (d, J=7.1 Hz, 1H, H2β), 5.44-5.21
 10 (m, 3H, NH, H3', H2'), 4.91 (d, J=9.9 Hz, 1H, H5), 4.62
 (m, 4H, methylenes), 4.44 (m, 1H, H7), 4.31 (d, J = 8.2
 Hz, 1H, H20α), 4.15 (d, J = 8.2 Hz, 1H, H20β), 3.90 (d,
 J=7.1 Hz, 1H, H3), 3.35 (d, J=5.5 Hz, 1H, 2'OH), 2.59 (m,
 1H, H6α), 2.49 (s, 3H, CH₃-imidazole), 2.36 (s, 3H, 4Ac),
 15 2.31 (m, 2H, H14), 2.13 (s, 3H, 10Ac), 1.95 (m, 1H, H6β),
 1.88 (br s, 3H, Me18), 1.77 (s, 3H, Me19), 1.70 (s, 1H,
 1OH), 1.34 (s, 9H, t-butyl), 1.23 (s, 3H, Me17), 1.15 (s,
 3H, Me16).

EXAMPLE 3

20 Preparation of N-debenzoyl-N-(t-butylcarbamoyl)-
 3'-desphenyl-3'-(4-nitrophenyl) taxol. (Taxoltere p-nip)



hn-2-66-3

To a solution of 7-triethylsilyl baccatin III
 25 (120 mg, 0.171 mmol) in 1.2 mL of THF at -45 °C was added

dropwise 0.104 mL of a 1.63 M solution of nBuLi in hexane. After 0.5 h at -45 °C, a solution of *cis*-1-(*t*-butoxycarbonyl)-3-triethylsilyloxy-4-(4-nitrophenyl)azetidin-2-one (361 mg, 0.885 mmol) in 1.2 mL of THF was added dropwise to the mixture. The solution was warmed to 0 °C and kept at that temperature for 1 h before 1 mL of a 10% solution of AcOH in THF was added. The mixture was partitioned between saturated aqueous NaHCO₃ and 60/40 ethyl acetate/hexane. Evaporation of the organic layer gave a residue which was purified by filtration through silica gel to give 192 mg of a mixture containing (2'R,3'S)-2',7-(bis)triethylsilyl-N-debenzoyl-N-(*t*-butylcarbamoyl)-3'-desphenyl-3'-(4-nitrophenyl) taxol and a very small amount of the (2'S,3'R) isomer.

To a solution of 192 mg of the mixture obtained from the previous reaction in 11 mL of acetonitrile and 0.55 mL of pyridine at 0 °C was added 1.7 mL of 48% aqueous HF. The mixture was stirred at 0 °C for 3 h, then at 25 °C for 13 h, and partitioned between saturated aqueous sodium bicarbonate and ethyl acetate. Evaporation of the ethyl acetate solution gave 153 mg of material which was purified by flash chromatography to give 140 mg (91%) of N-debenzoyl-N-(*t*-butylcarbamoyl)-3'-desphenyl-3'-(4-nitrophenyl) taxol, which was recrystallized from methanol/water.

m.p. 172-173 °C; $[\alpha]_{\text{Na}}^{25}$ -54 °C (c 0.0046, CHCl₃).

EXAMPLE 4

Biological Studies of Taxoltere Metro and Taxoltere p-nip

4.1. In vitro chemotherapeutic activity.

Chinese hamster ovary (CHO) and human colon carcinoma (HCT-116) cells were treated with different concentrations of taxol, taxoltere metro, or taxoltere p-nip at 37 °C for 24 hours. Cell survival was evaluated by the colony forming assay. As shown in Figures 1 and

2, both taxoltere metro and taxoltere p-nip are much more efficient than taxol in killing CHO and HCT-116 cells. At the 50% survival level, taxoltere metro is about 15 times, and taxoltere p-nip is about 45 times, more effective than taxol in killing CHO cells; measured at the 1% survival level, taxoltere metro is about 10 times, and taxoltere p-nip is about 30 times, more effective than taxol in killing HCT-116 cells. As illustrated in Figures 1 and 2, both taxoltere metro and taxoltere p-nip exhibit significantly stronger ability than taxol to kill both types of cells at every drug dose point.

4.2. **In vitro chemotherapeutic radiosensitization.** These studies were carried out as above except that cells were irradiated (General Electric Maxitron 300 at 250 kvp and 20 mA (HVL 20 mm Al filter; dose rate of 2 Gy/min) after two hours incubation. Figures 3 and 4 show the results of experiments in which cells were subjected to different radiation doses in the presence or absence of drugs. Both taxoltere metro and taxoltere p-nip strongly radiosensitize both CHO and HCT-116 cells, although taxol does not. For CHO cells, the sensitizer enhancement ratio (SER) is 2.3 for 100 nM taxoltere metro and 1.6 for taxoltere p-nip. For HCT-116 cells, taxoltere metro shows a SER of 1.2 at the (low) dose of 10 nM, and taxoltere p-nip has a SER of 1.5. At each radiation dose point, there is a significantly enhanced decrease in the surviving fraction for the taxoltere metro and taxoltere p-nip treated groups, but not for the taxol treated groups. HCT-116 cells are more sensitive to both taxoltere metro and taxoltere p-nip than CHO cells, hence lower concentrations of the drugs are required for a significant enhancement of radiation induced cell killing.

4.3. **In vivo drug toxicity.** Acute toxicity experiments were conducted on C3H/HeJ mice. The LD₅₀ (lethal dose to 50% of animals) values were determined by standard procedures described by Chan and Hayes (Chan, P.K. and Hayes, A.W. Principles and Methods for acute toxicity and eye irritancy. In Hayes, A.W., ed. Principles and Methods of Toxicity. 2nd Ed., New York, New York, Raven Press; 1989: 169220). The LD_{50/5} for i.p. taxoltere metro is 249.67 mg/kg, compared with 140.97 mg/kg for i.p. taxol. At high drug dose levels, the death of mice in the taxol treated groups occurred sooner than the death of mice in the taxoltere metro treated groups. Severe toxic symptoms such as the constriction of pupils and the contraction of erectile tissue of hair follicles (resulting in rough hair) were observed 24 hours after drug administration in the taxol treated groups, but not in the taxoltere metro treated groups. Obviously, the acute toxicity of taxoltere metro is significantly lower than that of taxol. The LD_{50/5} values are 79.13 mg/kg for i.p. administration of taxoltere p-nip and 134.16 mg/kg for i.v. administration of taxoltere p-nip. The data strongly demonstrate that taxoltere p-nip is significantly less toxic when administered i.v. than it is when administered i.p.

4.4. **In vivo chemotherapeutic activity.** Taxol, taxoltere metro, and taxoltere p-nip were administered at equitoxic doses using C3H/HeJ mice bearing a mouse mammary adenocarcinoma (MTG-B) in the right flank. As illustrated in Figure 5, taxol-metro is much more effective than taxol in increasing the life span of mammary tumor-bearing mice. The survival time for control mice was 13 days. At 40% of the LD_{50/5} dose, taxol increased the survival time by 31% to 17 days, and taxoltere metro increased the survival time by 123% to 29 days. Taxoltere p-nip (Figure 6) is substantially more

potent, increasing the life span by 1015% to 145 days (dose-response data not shown because only a single dose (40% of $LD_{50/5}$) was studied). Thus, under equitoxic conditions, taxoltere metro was 4 times, and taxoltere p-nip was 33 times, more effective than taxol.

Two experiments in which fractionated lower doses (20% of $LD_{50/5}$) of taxol, taxoltere metro and taxoltere p-nip were administered to MTG-B tumor bearing C3H/Hej mice have been carried out. In the first of these, drugs were administered on a q7d x 4 schedule (Figure 7). As shown in Figure 8, tumor doubling times are greatly extended by taxoltere metro and are even further extended by taxoltere p-nip on this administration schedule. In the second experiment (Figure 9), drugs were administered on a q1ld x4 schedule. Using this schedule, tumor doubling times (shown in Figure 10) are again greatly extended, with taxoltere metro producing the better results. Efficacy is positively correlated with frequency of injection (i.e., smaller intervals between injections). On both schedules, taxoltere metro and taxoltere p-nip exhibit significantly stronger antitumor effects than taxol.

4.5. In vivo chemotherapeutic radiosensitization. MTG-B tumor bearing C3H/Hej mice received single equitoxic drug doses i.p. (40% of $LD_{50/5}$). One day later the mice were placed in a lead holder (with the tumor-bearing hind leg exposed), and the tumor was subjected to a 22 Gy Xray dose (Figure 11). As shown in Figure 12, tumor doubling times (TDT) of 3.5 days (control), 8.5 days (22 Gy Xray exposure alone), 14.5 days (taxol alone), 19.5 days (taxol plus 22 Gy Xray), 20.5 days (taxoltere metro alone), and 29.5 days (taxoltere metro plus 22 Gy Xray) were observed. Equally large effects were observed when mice were given taxoltere metro two hours prior to irradiation (data not

shown). Taxoltere p-nip alone (TDT 3D 36.5 days) is even more effective in delaying tumor growth (Figure 12), and the combination of taxoltere p-nip plus 22 Gy produced a TDT of 65 days.

Administration of taxoltere p-nip i.v. (24% of $LD_{50/5}$) (Figure 13), gave a tumor doubling time of 52 days (Figure 14), approximately equivalent to the results obtained from i.p. administration.

4.6. **Cure Rates.** The term "cure" is defined by the U.S. National Cancer Institute as tumor-free survival for at least twice the survival time of control tumor-bearing mice, therefore we have used 28 days tumor-free to define a "cure" in this system. The cure rate is 40% for mice treated i.p. with taxoltere p-nip alone at a single dose, and 75% for the combination of taxoltere p-nip and 22 Gy (Figure 15). Similarly, a single i.v. injection of taxoltere p-nip alone induces a cure rate of 17%, and an 83% cure rate was observed for the combination of taxoltere p-nip and 22 Gy (Figure 16), even though this experiment was conducted with a lower dose (24% of $LD_{50/5}$). Although in this latter group one tumor recurred after 40 days, some treated mice have remained tumor-free for more than one year, and still survive.

As shown in Figure 17, administration of taxoltere p-nip iv at 40% of $LD_{50/5}$ produced a cure rate of 50%, and iv administration of taxoltere p-nip at 40% of $LD_{50/5}$ in combination with 22 Gy radiation produced a cure rate of 100%. Cure rate as a function of drug dose is shown in Figure 18.

In view of the above, it will be seen that the several objects of the invention are achieved and other advantageous results attained.

As various changes could be made in the above compounds and methods without departing from the scope of

the invention it is intended that all matter contained in the above description shall be interpreted as illustrative and not in a limiting sense.


WHAT IS CLAIMED IS:

1. A compound comprising a taxane containing at least 2 electron-affinic radiosensitizing functional groups.

2. The compound of claim 1 wherein at least one of the radiosensitizing groups is a nitro-substituted carbocyclic or heterocyclic aromatic moiety which is attached to the C2, C4, C7, C9, C10 or C14 position of the taxane.

3. The compound of claim 1 wherein said radiosensitizing groups are independently selected from nitro-substituted carbocyclic and heterocyclic aromatic moieties and wherein at least one of said radiosensitizing groups is attached to the C2, C4, C7, C9, C10 or C14 position of the taxane.

4. A compound corresponding to the structure:




wherein

- M comprises ammonium or is a metal;
- R₁ is hydrogen or hydroxy;
- R₂ is -OT₂, -OCOZ₂, -OCOOZ₂, RSG₁ or RSG₂;
- R₄ is -OT₄, -OCOZ₄, -OCOOZ₄, RSG₁ or RSG₂;
- R₇ is hydrogen, halogen, -OT₇, -OCOZ₇, -OCOOZ₇, RSG₁ or RSG₂;
- R₉ is hydrogen, keto, -OT₉, -OCOZ₉, -OCOOZ₉, RSG₁ or RSG₂;

R_{10} is hydrogen, keto, $-OT_{10}$, $-OCOZ_{10}$, $-OCOOZ_{10}$, RSG_1 or RSG_2 ;

R_7 , R_9 , and R_{10} independently have the alpha or
15 beta stereochemical configuration;

R_{13} is hydroxy, protected hydroxy, keto, $MO-$ or



R_{14} is hydrogen, hydroxy, protected hydroxy, RSG_1 or RSG_2 ;

20 T_2 , T_4 , T_7 , T_9 and T_{10} are independently hydrogen or hydroxy protecting group;

X_1 is $-OX_6$;

X_2 is hydrogen, hydrocarbon, heterosubstituted hydrocarbon, heteroaryl, or heterosubstituted heteroaryl;

25 X_3 is alkyl, heterosubstituted alkyl, alkenyl, heterosubstituted alkenyl, alkynyl, heterosubstituted alkynyl, phenyl, heteroaryl, or heterosubstituted heteroaryl;

30 X_4 is hydrogen, alkyl, heterosubstituted alkyl, alkenyl, heterosubstituted alkenyl, alkynyl, heterosubstituted alkynyl, phenyl, heteroaryl, or heterosubstituted heteroaryl;

X_5 is $-X_{10}$, $-OX_{10}$, $-SX_{11}$, or $-NX_8X_{11}$;

35 X_6 is hydrogen, hydrocarbon, heterosubstituted hydrocarbon, heteroaryl, heterosubstituted heteroaryl, hydroxy protecting group or a functional group which increases the water solubility of the taxane derivative;

X_8 is hydrogen, hydrocarbon, heterosubstituted hydrocarbon, RSG_1 or RSG_2 ;

40 X_{10} is alkyl, heterosubstituted alkyl, alkenyl, heterosubstituted alkenyl, alkynyl, heterosubstituted

alkynyl, phenyl, heteroaryl, or heterosubstituted heteroaryl;

45 X_{11} is hydrocarbon, heterosubstituted hydrocarbon, heteroaryl, heterosubstituted heteroaryl, RSG_1 or RSG_2 ;

Z_2 , Z_4 , Z_7 , Z_9 and Z_{10} are independently hydrocarbon, heterosubstituted hydrocarbon, heteroaryl, or heterosubstituted heteroaryl;

50 RSG_1 is an electron-affinic moiety;

RSG_2 is $-L-(RSG_1)_n$;

L is a linker comprising a chain of 1 to 30 atoms in the chain, the atoms being selected from the group consisting of C, O, N, S, Si, and P; and

55 n is an integer greater than or equal to 1; provided that the compound contains at least one radiosensitizing group attached to the C2, C4, C7, C9, C10, C14, C3' or C5' position of the compound.

5. The compound of claim 4 wherein RSG_1 is an electron-affinic group selected from the group consisting of (i) carbocyclic and heterocyclic aromatic moieties which possess one or more carbonyl, trifluoromethyl, halogen, nitro, sulfonyl, sulfinyl, phosphoryl, oxide or cyano groups, (ii) heterocyclic aromatic moieties containing two or more heteroatoms, (iii) metal complexes, and (iv) organo-metallic groups in which the metal is covalently bonded to carbon.

6. The compound of claim 4 wherein RSG_1 is selected from the group consisting of imidazoles, triazoles, pyridines, benzamides, nicotinamides, benzotriazine oxides, furans, thiophenes, oxazoles and thiozoles possessing one or more carbonyl, trifluoromethyl, halogen, nitro, sulfonyl, sulfinyl, phosphoryl, oxide or cyano groups.

50

7. The compound of claim 4 wherein

R_1 is hydrogen or hydroxy;

R_2 is $-\text{OCOZ}_2$, RSG_1 , or RSG_2 ;


R_4 is $-\text{OCOZ}_4$, RSG_1 or RSG_2 ;

5 R_7 is hydrogen, halogen, $-\text{OT}_7$, $-\text{OCOZ}_7$, RSG_1 or RSG_2 ;

R_9 is hydrogen, keto, $-\text{OT}_9$, $-\text{OCOZ}_9$, RSG_1 or RSG_2 ;

R_{10} is hydrogen, keto, $-\text{OT}_{10}$, $-\text{OCOZ}_{10}$, RSG_1 or RSG_2 ;

10 R_{13} is



R_{14} is hydrogen, hydroxy or protected hydroxy;

T_2 , T_4 , T_7 , T_9 and T_{10} are independently hydrogen or hydroxy protecting group;

15 X_1 is $-\text{OX}_6$;

X_2 is hydrogen;

X_3 is alkyl, alkenyl, phenyl, heteroaryl, or heterosubstituted heteroaryl;

20 X_4 is hydrogen, hydrocarbon, heteroaryl, or heterosubstituted heteroaryl;

X_5 is $-\text{X}_{10}$, $-\text{OX}_{10}$, $-\text{SX}_{11}$, or $-\text{NX}_8\text{X}_{11}$;

X_6 is hydrogen or hydroxy protecting group;

X_8 is hydrogen, hydrocarbon, heterosubstituted hydrocarbon, RSG_1 or RSG_2 ;

25 X_{10} is alkyl, alkenyl, phenyl, heteroaryl, or heterosubstituted heteroaryl;

Z_2 , Z_4 , Z_7 , Z_9 and Z_{10} are independently hydrocarbon, heterosubstituted hydrocarbon, heteroaryl, or heterosubstituted heteroaryl;

30 RSG_1 is an electron-affinic moiety;

RSG_2 is $-\text{L}-(\text{RSG}_1)_n$;


L is a linker comprising a chain of 1 to 10 atoms in the chain, the atoms being selected from the group consisting of C, O, N, S, Si, and P; and
 35 n is 1 or 2.

8. The compound of claim 4 wherein RSG₁ is a heterocyclic aromatic moiety containing two or more heteroatoms.

9. The compound of claim 4 wherein RSG₁ is a metal complex.

10. The compound of claim 4 wherein RSG₁ is selected from the group consisting imidazoles, triazoles, pyridines, benzamides, furans, thiophenes, oxazoles and thiozoles possessing one or more nitro groups.

11. A compound corresponding to the structure:




wherein

- 5 M comprises ammonium or is a metal;
 R₁ is hydrogen or hydroxy;
 R₂ is RSG₁ or RSG₂;
 R₄ is -OT₄, -OCOOZ₄, -OCOOZ₄, RSG₁ or RSG₂;
 R₇ is hydrogen, halogen, -OT₇, -OCOOZ₇, -OCOOZ₇,
 RSG₁ or RSG₂;
 10 R₉ is hydrogen, keto, -OT₉, -OCOOZ₉, -OCOOZ₉, RSG₁
 or RSG₂;
 R₁₀ is hydrogen, keto, -OT₁₀, -OCOOZ₁₀, -OCOOZ₁₀,
 RSG₁ or RSG₂;

R_7 , R_9 , and R_{10} independently have the alpha or
 15 beta stereochemical configuration;

R_{13} is hydroxy, protected hydroxy, keto, MO- or



R_{14} is hydrogen, hydroxy, protected hydroxy,
 RSG₁ or RSG₂;

20 T_2 , T_4 , T_7 , T_9 and T_{10} are independently hydrogen
 or hydroxy protecting group;

X_1 is $-OX_6$;

X_2 is hydrogen, hydrocarbon, heterosubstituted
 hydrocarbon, heteroaryl, or heterosubstituted heteroaryl;

25 X_3 and X_4 are independently hydrogen,
 hydrocarbon, heterosubstituted hydrocarbon, heteroaryl,
 heterosubstituted heteroaryl or RSG₁;

X_5 is $-X_{10}$, $-OX_{10}$, $-SX_{10}$, or $-NX_8X_{10}$;

X_6 is hydrogen, hydrocarbon, heterosubstituted
 30 hydrocarbon, heteroaryl, heterosubstituted heteroaryl,
 hydroxy protecting group or a functional group which
 increases the water solubility of the taxane derivative;

X_8 is hydrogen, hydrocarbon, heterosubstituted
 hydrocarbon, RSG₁ or RSG₂;

35 X_{10} is hydrocarbon, heterosubstituted
 hydrocarbon, heteroaryl, heterosubstituted heteroaryl,
 RSG₁ or RSG₂;


Z_4 , Z_7 , Z_9 and Z_{10} are independently hydrocarbon,
 heterosubstituted hydrocarbon, heteroaryl, or
 40 heterosubstituted heteroaryl;

RSG₁ is an electron-affinic moiety;

RSG₂ is $-L-(RSG_1)_n$;


L is a linker comprising a chain of 1 to 30 atoms in the chain, the atoms being selected from the group consisting of C, O, N, S, Si, and P; and
 45 n is an integer greater than or equal to 1.

12. A compound corresponding to the structure:



wherein

- M comprises ammonium or is a metal;
 5 R₁ is hydrogen or hydroxy;
 R₂ is -OT₂, -OCOOZ₂, -OCOOZ₂, RSG₁ or RSG₂;
 R₄ is RSG₁ or RSG₂;
 R₇ is hydrogen, halogen, -OT₇, -OCOOZ₇, -OCOOZ₇,
 RSG₁ or RSG₂;
 10 R₉ is hydrogen, keto, -OT₉, -OCOOZ₉, -OCOOZ₉, RSG₁
 or RSG₂;
 R₁₀ is hydrogen, keto, -OT₁₀, -OCOOZ₁₀, -OCOOZ₁₀,
 RSG₁ or RSG₂;
 R₇, R₉, and R₁₀ independently have the alpha or
 15 beta stereochemical configuration;
 R₁₃ is hydroxy, protected hydroxy, keto, MO- or




R₁₄ is hydrogen, hydroxy, protected hydroxy,
 RSG₁ or RSG₂;

- 20 T_2 , T_4 , T_7 , T_9 and T_{10} are independently hydrogen or hydroxy protecting group;
- X_1 is $-OX_6$;
- X_2 is hydrogen, hydrocarbon, heterosubstituted hydrocarbon, heteroaryl, or heterosubstituted heteroaryl;
- 25 X_3 and X_4 are independently hydrogen, hydrocarbon, heterosubstituted hydrocarbon, heteroaryl, heterosubstituted heteroaryl, or RSG_1 ;
- X_5 is $-X_{10}$, $-OX_{10}$, $-SX_{10}$, or $-NX_8X_{10}$;
- X_6 is hydrogen, hydrocarbon, heterosubstituted hydrocarbon, heteroaryl, heterosubstituted heteroaryl,
- 30 hydroxy protecting group or a functional group which increases the water solubility of the taxane derivative;
- X_8 is hydrogen, hydrocarbon, heterosubstituted hydrocarbon, RSG_1 or RSG_2 ;
- 35 X_{10} is hydrocarbon, heterosubstituted hydrocarbon, heteroaryl, heterosubstituted heteroaryl, RSG_1 or RSG_2 ;
- Z_2 , Z_7 , Z_9 and Z_{10} are independently hydrocarbon, heterosubstituted hydrocarbon, heteroaryl, or
- 40 heterosubstituted heteroaryl;
- RSG_1 is an electron-affinic moiety;
- RSG_2 is $-L-(RSG_1)_n$;
- L is a linker comprising a chain of 1 to 30 atoms in the chain, the atoms being selected from the
- 45 group consisting of C, O, N, S, Si, and P; and
- n is an integer greater than or equal to 1.

55

13. A compound corresponding to the structure:



wherein

M comprises ammonium or is a metal;

5

R_1 is hydrogen or hydroxy;

R_2 is $-OT_2$, $-OCOZ_2$, $-OCOOZ_2$, RSG_1 or RSG_2 ;

R_4 is $-OT_4$, $-OCOZ_4$, $-OCOOZ_4$, RSG_1 or RSG_2 ;


R_7 is RSG_1 or RSG_2 ;

10 R_9 is hydrogen, keto, $-OT_9$, $-OCOZ_9$, $-OCOOZ_9$, RSG_1 or RSG_2 ;

R_{10} is hydrogen, keto, $-OT_{10}$, $-OCOZ_{10}$, $-OCOOZ_{10}$, RSG_1 or RSG_2 ;

R_7 , R_9 , and R_{10} independently have the alpha or beta stereochemical configuration;

15 R_{13} is hydroxy, protected hydroxy, keto, MO- or



;

R_{14} is hydrogen, hydroxy, protected hydroxy, RSG_1 or RSG_2 ;


20 T_2 , T_4 , T_9 and T_{10} are independently hydrogen or hydroxy protecting group;

X_1 is $-OX_6$;

X_2 is hydrogen, hydrocarbon, heterosubstituted hydrocarbon, heteroaryl, or heterosubstituted heteroaryl;

- X_3 and X_4 are independently hydrogen,
 25 hydrocarbon, heterosubstituted hydrocarbon, heteroaryl,
 heterosubstituted heteroaryl or RSG_1 ;
 X_5 is $-X_{10}$, $-OX_{10}$, $-SX_{10}$, or $-NX_8X_{10}$;
 X_6 is hydrogen, hydrocarbon, heterosubstituted
 hydrocarbon, heteroaryl, heterosubstituted heteroaryl,
 30 hydroxy protecting group or a functional group which
 increases the water solubility of the taxane derivative;
 X_8 is hydrogen, hydrocarbon, heterosubstituted
 hydrocarbon, RSG_1 or RSG_2 ;
 X_{10} is hydrocarbon, heterosubstituted
 35 hydrocarbon, heteroaryl, heterosubstituted heteroaryl,
 RSG_1 or RSG_2 ;
 Z_2 , Z_4 , Z_9 and Z_{10} are independently hydrocarbon,
 heterosubstituted hydrocarbon, heteroaryl, or
 heterosubstituted heteroaryl;
 40 RSG_1 is an electron-affinic moiety;
 RSG_2 is $-L-(RSG_1)_n$;
 L is a linker comprising a chain of 1 to 30
 atoms in the chain, the atoms being selected from the
 group consisting of C, O, N, S, Si, and P; and
 45 n is an integer greater than or equal to 1.

14. A compound corresponding to the structure:



wherein

- M comprises ammonium or is a metal;
 5 R_1 is hydrogen or hydroxy;
 R_2 is $-OT_2$, $-OCOZ_2$, $-OCOOZ_2$, RSG_1 or RSG_2 ;
 R_4 is $-OT_4$, $-OCOZ_4$, $-OCOOZ_4$, RSG_1 or RSG_2 ;


R_7 is hydrogen, keto, $-OT_7$, $-OCOZ_7$, $-OCOOZ_7$, RSG_1 or RSG_2 ;

10 R_9 is RSG_1 or RSG_2 ;

R_{10} is hydrogen, keto, $-OT_{10}$, $-OCOZ_{10}$, $-OCOOZ_{10}$, RSG_1 or RSG_2 ;

R_7 , R_9 , and R_{10} independently have the alpha or beta stereochemical configuration;

15 R_{13} is hydroxy, protected hydroxy, keto, MO- or



R_{14} is hydrogen, hydroxy, protected hydroxy, RSG_1 or RSG_2 ;

20 T_2 , T_4 , T_7 and T_{10} are independently hydrogen or hydroxy protecting group;

X_1 is $-OX_6$;

X_2 is hydrogen, hydrocarbon, heterosubstituted hydrocarbon, heteroaryl, or heterosubstituted heteroaryl;

25 X_3 and X_4 are independently hydrogen, hydrocarbon, heterosubstituted hydrocarbon, heteroaryl, heterosubstituted heteroaryl or RSG_1 ;

X_5 is $-X_{10}$, $-OX_{10}$, $-SX_{10}$, or $-NX_8X_{10}$;

30 X_6 is hydrogen, hydrocarbon, heterosubstituted hydrocarbon, heteroaryl, heterosubstituted heteroaryl, hydroxy protecting group or a functional group which increases the water solubility of the taxane derivative;

X_8 is hydrogen, hydrocarbon, heterosubstituted hydrocarbon, RSG_1 or RSG_2 ;

35 X_{10} is hydrocarbon, heterosubstituted hydrocarbon, heteroaryl, heterosubstituted heteroaryl, RSG_1 or RSG_2 ;

58

Z_2 , Z_4 , Z_7 , and Z_{10} are independently hydrocarbon, heterosubstituted hydrocarbon, heteroaryl, or heterosubstituted heteroaryl;


40 RSG_1 is an electron-affinic moiety;

RSG_2 is $-L-(RSG_1)_n$;

L is a linker comprising a chain of 1 to 30 atoms in the chain, the atoms being selected from the group consisting of C, O, N, S, Si, and P; and

45 n is an integer greater than or equal to 1.

15. A compound corresponding to the structure:



wherein

M comprises ammonium or is a metal;

5 R_1 is hydrogen or hydroxy;

R_2 is $-OT_2$, $-OCOT_2$, $-OCOT_2$, RSG_1 or RSG_2 ;

R_4 is $-OT_4$, $-OCOT_4$, $-OCOT_4$, RSG_1 or RSG_2 ;


R_7 is hydrogen, keto, $-OT_7$, $-OCOT_7$, $-OCOT_7$, RSG_1 or RSG_2 ;

10 R_9 is hydrogen, keto, $-OT_9$, $-OCOT_9$, $-OCOT_9$, RSG_1 or RSG_2 ;

R_{10} is RSG_1 or RSG_2 ;

R_7 , R_9 , and R_{10} independently have the alpha or beta stereochemical configuration;

15 R_{13} is hydroxy, protected hydroxy, keto, MO- or



R_{14} is hydrogen, hydroxy, protected hydroxy, RSG₁ or RSG₂;

20 T_2 , T_4 , T_7 and T_9 are independently hydrogen or hydroxy protecting group;

X_1 is -OX₆;

X_2 is hydrogen, hydrocarbon, heterosubstituted hydrocarbon, heteroaryl, or heterosubstituted heteroaryl;

25 X_3 and X_4 are independently hydrogen, hydrocarbon, heterosubstituted hydrocarbon, heteroaryl, heterosubstituted heteroaryl or RSG₁;

X_5 is -X₁₀, -OX₁₀, -SX₁₀, or -NX₉X₁₀;

30 X_6 is hydrogen, hydrocarbon, heterosubstituted hydrocarbon, heteroaryl, heterosubstituted heteroaryl, hydroxy protecting group or a functional group which increases the water solubility of the taxane derivative;

X_8 is hydrogen, hydrocarbon, heterosubstituted hydrocarbon, RSG₁ or RSG₂;

35 X_{10} is hydrocarbon, heterosubstituted hydrocarbon, heteroaryl, heterosubstituted heteroaryl, RSG₁ or RSG₂;

Z_2 , Z_4 , Z_7 and Z_9 are independently hydrocarbon, heterosubstituted hydrocarbon, heteroaryl, or heterosubstituted heteroaryl;


40 RSG₁ is an electron-affinic moiety;

RSG₂ is -L-(RSG₁)_n;

L is a linker comprising a chain of 1 to 30 atoms in the chain, the atoms being selected from the group consisting of C, O, N, S, Si, and P; and

45 n is an integer greater than or equal to 1.

16. A compound corresponding to the structure:



wherein

M comprises ammonium or is a metal;

5 R_1 is hydrogen or hydroxy;

R_2 is $-OT_2$, $-OC(O)Z_2$, $-OC(O)OZ_2$, RSG_1 or RSG_2 ;

R_4 is $-OT_4$, $-OC(O)Z_4$, $-OC(O)OZ_4$, RSG_1 or RSG_2 ;


R_7 is hydrogen, keto, $-OT_7$, $-OC(O)Z_7$, $-OC(O)OZ_7$, RSG_1 or RSG_2 ;

10 R_9 is hydrogen, keto, $-OT_9$, $-OC(O)Z_9$, $-OC(O)OZ_9$, RSG_1 or RSG_2 ;

R_{10} is hydrogen, keto, $-OT_{10}$, $-OC(O)Z_{10}$, $-OC(O)OZ_{10}$, RSG_1 or RSG_2 ;

15 R_7 , R_9 , and R_{10} independently have the alpha or beta stereochemical configuration;

R_{13} is hydroxy, protected hydroxy, keto, MO- or



R_{14} is RSG_1 or RSG_2 ;

20 T_2 , T_4 , T_7 , T_9 and T_{10} are independently hydrogen or hydroxy protecting group;

X_1 is $-OX_6$;

X_2 is hydrogen, hydrocarbon, heterosubstituted hydrocarbon, heteroaryl, or heterosubstituted heteroaryl;

25 X_3 and X_4 are independently hydrogen, hydrocarbon, heterosubstituted hydrocarbon, heteroaryl, heterosubstituted heteroaryl or RSG_1 ;

X_5 is $-X_{10}$, $-OX_{10}$, $-SX_{10}$, or $-NX_8X_{10}$;

30 X_6 is hydrogen, hydrocarbon, heterosubstituted hydrocarbon, heteroaryl, heterosubstituted heteroaryl, hydroxy protecting group or a functional group which increases the water solubility of the taxane derivative;

X_8 is hydrogen, hydrocarbon, heterosubstituted hydrocarbon, RSG_1 or RSG_2 ;

35 X_{10} is hydrocarbon, heterosubstituted hydrocarbon, heteroaryl, heterosubstituted heteroaryl, RSG_1 or RSG_2 ;

Z_2 , Z_4 , Z_7 , Z_9 and Z_{10} are independently hydrocarbon, heterosubstituted hydrocarbon, heteroaryl or heterosubstituted heteroaryl;

40 RSG_1 is an electron-affinic moiety;

RSG_2 is $-L-(RSG_1)_n$;

L is a linker comprising a chain of 1 to 30 atoms in the chain, the atoms being selected from the group consisting of C, O, N, S, Si, and P; and

45 n is an integer greater than or equal to 1.

17. A method of killing tumor cells in a warm-blooded animal, the method comprising:

5 (a) administering to the warm-blooded animal a taxane containing an electron-affinic radiosensitizing functional group,

(b) followed by, after a time interval sufficient to enhance radiosensitization of the tumor cells, irradiating the tumor cells with a dose of radiation effective to kill the tumor cells.

18. A method as set forth in claim 16 further comprising heat treating the tumor cells.

AMENDED CLAIMS


[received by the International Bureau on 4 February 1997((04.02.97); original claims 4 and 11-16 amended; remaining claims unchanged (7 pages)]

1. A compound comprising a taxane containing at least 2 electron-affinic radiosensitizing functional groups.

2. The compound of claim 1 wherein at least one of the radiosensitizing groups is a nitro-substituted carbocyclic or heterocyclic aromatic moiety which is attached to the C2, C4, C7, C9, C10 or C14 position of the taxane.

3. The compound of claim 1 wherein said radiosensitizing groups are independently selected from nitro-substituted carbocyclic and heterocyclic aromatic moieties and wherein at least one of said radiosensitizing groups is attached to the C2, C4, C7, C9, C10 or C14 position of the taxane.

4. A compound corresponding to the structure:



wherein

M comprises ammonium or is a metal;

5 R₁ is hydrogen or hydroxy;

R₂ is -OT₂, -OCOT₂, -OCOT₂, RSG₁ or RSG₂;

R₄ is -OT₄, -OCOT₄, -OCOT₄, RSG₁ or RSG₂;

R₇ is hydrogen, halogen, -OT₇, -OCOT₇, -OCOT₇, RSG₁ or RSG₂;

10 R₉ is hydrogen, keto, -OT₉, -OCOT₉, -OCOT₉, RSG₁ or RSG₂;

L is a linker comprising a chain of 1 to 10 atoms in the chain, the atoms being selected from the group consisting of C, O, N, S, Si, and P; and


35 n is 1 or 2.

8. The compound of claim 4 wherein RSG_1 is a heterocyclic aromatic moiety containing two or more heteroatoms.

9. The compound of claim 4 wherein RSG_1 is a metal complex.

10. The compound of claim 4 wherein RSG_1 is selected from the group consisting imidazoles, triazoles, pyridines, benzamides, furans, thiophenes, oxazoles and thiozoles possessing one or more nitro groups.

11. A compound corresponding to the structure:



wherein

M comprises ammonium or is a metal;

5 R_1 is hydrogen or hydroxy;

R_2 is RSG_1 or RSG_2 ;

R_4 is $-OT_4$, $-OCOT_4$, $-OCOT_4$, RSG_1 or RSG_2 ;


R_7 is hydrogen, halogen, $-OT_7$, $-OCOT_7$, $-OCOT_7$, RSG_1 or RSG_2 ;

10 R_9 is hydrogen, keto, $-OT_9$, $-OCOT_9$, $-OCOT_9$, RSG_1 or RSG_2 ;

R_{10} is hydrogen, keto, $-OT_{10}$, $-OCOT_{10}$, $-OCOT_{10}$, RSG_1 or RSG_2 ;


L is a linker comprising a chain of 1 to 30 atoms in
 45 the chain, the atoms being selected from the group
 consisting of C, O, N, S, Si, and P; and
 n is an integer greater than or equal to 1.

12. A compound corresponding to the structure:




wherein

- M comprises ammonium or is a metal;
- 5 R_1 is hydrogen or hydroxy;
- R_2 is $-OT_2$, $-OCOOZ_2$, $-OCOOZ_2$, RSG_1 or RSG_2 ;
- R_4 is RSG_1 or RSG_2 ;
- R_7 is hydrogen, halogen, $-OT_7$, $-OCOOZ_7$, $-OCOOZ_7$, RSG_1
 or RSG_2 ;
- 10 R_9 is hydrogen, keto, $-OT_9$, $-OCOOZ_9$, $-OCOOZ_9$, RSG_1 or
 RSG_2 ;
- R_{10} is hydrogen, keto, $-OT_{10}$, $-OCOOZ_{10}$, $-OCOOZ_{10}$, RSG_1 or
 RSG_2 ;
- R_7 , R_9 , and R_{10} independently have the alpha or beta
 15 stereochemical configuration;
- R_{13} is hydroxy, protected hydroxy, keto, MO- or




R_{14} is hydrogen, hydroxy, protected hydroxy, RSG_1 or
 RSG_2 ;

13. A compound corresponding to the structure:



wherein


- 5 M comprises ammonium or is a metal;
 R₁ is hydrogen or hydroxy;
 R₂ is -OT₂, -OCOZ₂, -OCOOZ₂, RSG₁ or RSG₂;
 R₄ is -OT₄, -OCOZ₄, -OCOOZ₄, RSG₁ or RSG₂;
 R₇ is RSG₁ or RSG₂;
 10 R₉ is hydrogen, keto, -OT₉, -OCOZ₉, -OCOOZ₉, RSG₁ or RSG₂;
 R₁₀ is hydrogen, keto, -OT₁₀, -OCOZ₁₀, -OCOOZ₁₀, RSG₁ or RSG₂;
 R₇, R₉, and R₁₀ independently have the alpha or beta stereochemical configuration;
 15 R₁₃ is hydroxy, protected hydroxy, keto, MO- or



- R₁₄ is hydrogen, hydroxy, protected hydroxy, RSG₁ or RSG₂;
 20 T₂, T₄, T₉ and T₁₀ are independently hydrogen or hydroxy protecting group;
 X₁ is -OX₆;
 X₂ is hydrogen, hydrocarbon, heterosubstituted hydrocarbon, heteroaryl, or heterosubstituted heteroaryl;

- 25 X_3 and X_4 are independently hydrogen, hydrocarbon, heterosubstituted hydrocarbon, heteroaryl, heterosubstituted heteroaryl or RSG_1 ;
 X_5 is $-X_{10}$, $-OX_{10}$, $-SX_{10}$, or $-NX_8X_{10}$;
 X_6 is hydrogen, hydrocarbon, heterosubstituted
30 hydrocarbon, heteroaryl, heterosubstituted heteroaryl, hydroxy protecting group or a functional group which increases the water solubility of the taxane derivative;
 X_8 is hydrogen, hydrocarbon, heterosubstituted hydrocarbon, RSG_1 or RSG_2 ;
35 X_{10} is hydrocarbon, heterosubstituted hydrocarbon, heteroaryl, heterosubstituted heteroaryl, RSG_1 or RSG_2 ;
 Z_2 , Z_4 , Z_9 and Z_{10} are independently hydrocarbon, heterosubstituted hydrocarbon, heteroaryl, or heterosubstituted heteroaryl;
40 RSG_1 is an electron-affinic moiety;
 RSG_2 is $-L-(RSG_1)_n$;
 L is a linker comprising a chain of 1 to 30 atoms in the chain, the atoms being selected from the group consisting of C, O, N, S, Si, and P; and
45 n is an integer greater than or equal to 1.

14. A compound corresponding to the structure:



wherein

- M comprises ammonium or is a metal;
5 R_1 is hydrogen or hydroxy;
 R_2 is $-OT_2$, $-OCOT_2$, $-OCOT_2$, RSG_1 or RSG_2 ;
 R_4 is $-OT_4$, $-OCOT_4$, $-OCOT_4$, RSG_1 or RSG_2 ;


Z_2 , Z_4 , Z_7 , and Z_{10} are independently hydrocarbon, heterosubstituted hydrocarbon, heteroaryl, or
 40 heterosubstituted heteroaryl;

RSG_1 is an electron-affinic moiety;

RSG_2 is $-L-(RSG_1)_n$;


L is a linker comprising a chain of 1 to 30 atoms in the chain, the atoms being selected from the
 45 group consisting of C, O, N, S, Si, and P; and
 n is an integer greater than or equal to 1.

15. A compound corresponding to the structure:




wherein

- M comprises ammonium or is a metal;
 5 R_1 is hydrogen or hydroxy;
 R_2 is $-OT_2$, $-OCOOZ_2$, $-OCOOZ_2$, RSG_1 or RSG_2 ;
 R_4 is $-OT_4$, $-OCOOZ_4$, $-OCOOZ_4$, RSG_1 or RSG_2 ;
 R_7 is hydrogen, keto, $-OT_7$, $-OCOOZ_7$, $-OCOOZ_7$, RSG_1
 or RSG_2 ;
 10 R_9 is hydrogen, keto, $-OT_9$, $-OCOOZ_9$, $-OCOOZ_9$, RSG_1
 or RSG_2 ;
 R_{10} is RSG_1 or RSG_2 ;
 R_7 , R_9 , and R_{10} independently have the alpha or
 beta stereochemical configuration;
 15 R_{13} is hydroxy, protected hydroxy, keto, $MO-$ or



16. A compound corresponding to the structure:



wherein

M comprises ammonium or is a metal;

5

R_1 is hydrogen or hydroxy;

R_2 is $-OT_2$, $-OCOOZ_2$, $-OCOOZ_2$, RSG_1 or RSG_2 ;

R_4 is $-OT_4$, $-OCOOZ_4$, $-OCOOZ_4$, RSG_1 or RSG_2 ;

R_7 is hydrogen, keto, $-OT_7$, $-OCOOZ_7$, $-OCOOZ_7$, RSG_1

or RSG_2 ;

10

R_9 is hydrogen, keto, $-OT_9$, $-OCOOZ_9$, $-OCOOZ_9$, RSG_1

or RSG_2 ;

R_{10} is hydrogen, keto, $-OT_{10}$, $-OCOOZ_{10}$, $-OCOOZ_{10}$,


RSG_1 or RSG_2 ;

R_7 , R_9 , and R_{10} independently have the alpha or

15

beta stereochemical configuration;

R_{13} is hydroxy, protected hydroxy, keto, MO- or



;

R_{14} is RSG_1 or RSG_2 ;

T_2 , T_4 , T_7 , T_9 and T_{10} are independently hydrogen

20

or hydroxy protecting group;

X_1 is $-OX_6$;

X_2 is hydrogen, hydrocarbon, heterosubstituted hydrocarbon, heteroaryl, or heterosubstituted heteroaryl;

1/18




FIG. 1




FIG. 2

3/18





FIG. 3

FIG. 4



5/18




FIG. 5




FIG. 6

7/18




FIG. 7

8/18




FIG. 8

9/18

FIG. 9





FIG. 10



11/18

FIG. 11




FIG. 12





FIG. 13



14/18




FIG. 14

FIG. 15






FIG. 16



17/18

FIG. 17



18/18




FIG. 18

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US96/14613

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : A61K 31/335; C07D 305/14

US CL : 514/449; 549/510, 511

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/449; 549/510, 511

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
NONE

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
none

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X, P	US 5,466,834 A (HOLTON) 14 November 1995, see the entire document.	1-18



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:	* T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
* A* document defining the general state of the art which is not considered to be of particular relevance	* X*	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
* E* earlier document published on or after the international filing date	* Y*	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
* L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	* &*	document member of the same patent family
* O* document referring to an oral disclosure, use, exhibition or other means		
* P* document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

22 NOVEMBER 1996

Date of mailing of the international search report

05 DEC 1996

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

BA K. TRINH

Telephone No. (703) 308-1235



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification⁶: A61K 31/335, C07D 305/14</p>	<p>A1</p>	<p>(11) International Publication Number: WO 97/09979 (43) International Publication Date: 20 March 1997 (20.03.97)</p>
<p>(21) International Application Number: PCT/US96/14613 (22) International Filing Date: 13 September 1996 (13.09.96) (30) Priority Data: 60/003,687 13 September 1995 (13.09.95) US (71) Applicant (for all designated States except US): FLORIDA STATE UNIVERSITY [US/US]; Sliger Building, Room 100, 2035 East Paul Dirac Drive, Tallahassee, FL 32310 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): HOLTON, Robert, A. [US/US]; Dittmer Laboratory of Chemistry, Chemistry Dept., Room 715, Tallahassee, FL 32306 (US). NADIZADEH, Hossain [IR/US]; 415 Chapel Drive #D308, Tallahassee, FL 32304 (US). HOFER, Kurt, G. [US/US]; 235 Westridge Drive, Tallahassee, FL 32304 (US). YANG, Li-Xi [CN/US]; Apartment 4B, 1839 Miccosukee Road, Tallahassee, FL 32304 (US). (74) Agents: HEJLEK, Edward, J. et al.; Senniger, Powers, Leavitt and Roedel, 16th floor, One Metropolitan Square, St. Louis, MO 63102 (US).</p>		<p>(81) Designated States: AU, CA, CN, CZ, FI, HU, JP, KR, NO, NZ, PL, RU, US, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published With international search report. With amended claims.</p>
<p>(54) Title: RADIOSENSITIZING TAXANES AND THEIR PHARMACEUTICAL PREPARATIONS (57) Abstract Taxanes containing electron-affinic radiosensitizing functional groups, their pharmaceutical preparations, and methods of making and using this new class of highly potent radiosensitizers of tumor cells.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/335, C07D 305/14	A1	(11) International Publication Number: WO 97/09979 (43) International Publication Date: 20 March 1997 (20.03.97)
(21) International Application Number: PCT/US96/14613 (22) International Filing Date: 13 September 1996 (13.09.96) (30) Priority Data: 60/003,687 13 September 1995 (13.09.95) US (71) Applicant (for all designated States except US): FLORIDA STATE UNIVERSITY [US/US]; Sliger Building, Room 100, 2035 East Paul Dirac Drive, Tallahassee, FL 32310 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): HOLTON, Robert, A. [US/US]; Dittmer Laboratory of Chemistry, Chem- istry Dept., Room 715, Tallahassee, FL 32306 (US). NADIZADEH, Hossain [IR/US]; 415 Chapel Drive #D308, Tallahassee, FL 32304 (US). HOFER, Kurt, G. [US/US]; 235 Westridge Drive, Tallahassee, FL 32304 (US). YANG, Li-Xi [CN/US]; Apartment 4B, 1839 Miccosukee Road, Tallahassee, FL 32304 (US). (74) Agents: HEJLEK, Edward, J. et al.; Senniger, Powers, Leavitt and Roedel, 16th floor, One Metropolitan Square, St. Louis, MO 63102 (US).		(81) Designated States: AU, CA, CN, CZ, FI, HU, JP, KR, NO, NZ, PL, RU, US, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i> <i>With amended claims.</i>
(54) Title: RADIOSENSITIZING TAXANES AND THEIR PHARMACEUTICAL PREPARATIONS (57) Abstract <p>Taxanes containing electron-affinic radiosensitizing functional groups, their pharmaceutical preparations, and methods of making and using this new class of highly potent radiosensitizers of tumor cells.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	"	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

**RADIOSENSITIZING TAXANES AND
THEIR PHARMACEUTICAL PREPARATIONS**

Background of the Invention

This invention relates to novel
5 radiosensitizing compounds, and in particular, to
substituted taxanes containing radiosensitizing moieties,
their pharmaceutical preparations, and methods of using
this new class of highly potent radiosensitizers of tumor
cells.

10 In the United States alone, over a half million
patients undergo radiation therapy each year as a part of
their battle against cancer. To date, however, radiation
therapy has produced only limited success as a cancer
treatment. Understandably, therefore, a major effort has
15 been underway for a number of years to develop means to
improve the efficacy of such radiotherapy techniques.

It is widely believed that the presence of
radioresistant, hypoxic (poorly oxygenated) cells in
tumors constitutes a significant factor in causing local
20 failure in conventional cancer radiotherapy. For
example, it was reported by Gatenby et al., Int. J.
Radiat. Oncol. Biol. Phys. 14: 831-833 (1988), that for
head and neck tumors, the hypoxic cell volume is
inversely correlated with tumor radiosensitivity. Other
25 reports confirm this conclusion for a variety of types of
tumors and suggest that the presence of a concentration
of as little as 2-3% hypoxic cells in a tumor may double
the radiation dose required for tumor control.

Various solutions have been proposed to
30 overcome the problem of hypoxia, including carrying out
radiation treatments in high pressure oxygen chambers and
the substitution of "fast neutron" or π meson radiation
in place of X-rays. However, these techniques are not
wholly satisfactory for a number of reasons, including

the great expense and difficulty frequently associated with such procedures.

One promising field of investigation for dealing with radioresistant hypoxic tumor cells has been the use of "radiosensitizing" compounds which selectively increase the sensitivity of hypoxic cells to radiation. This specificity to hypoxic cells is also valuable because a significant percentage of solid tumors are characterized by such cells while most normal tissue is not. Thus, treatment with such compounds serves to enhance the impact of radiation on tumor cells while having little effect on the impact of radiation on healthy cell tissue. A number of heterocyclic, electron-affinic compounds, and in particular, those with oxidized nitrogen moieties, have been successfully used for the purpose of radiosensitizing hypoxic tumor cells. Specifically, the discovery that the nitroimidazoles, metronidazole (metro) and misonidazole (miso), sensitize hypoxic cells to radiation provided initial optimism for a breakthrough solution to the problem of tumor hypoxia. Unfortunately, however, both agents have proven to be highly toxic at therapeutic levels.

The possibility of using chemotherapeutic agents to selectively enhance radiation response in tumors has also been proposed. In addition to its use as a chemotherapeutic agent, taxol, for example, has been investigated in vitro and in vivo as a potential radiosensitizing drug. See, Tishler et al., Radiation Oncology Biol. Phys., 22:613-617 (1992); Tishler et al., Cancer Research, 52:3495-3497 (1992); Steren, et al., Gynecologic Oncology, 48:252-258 (1993); Steren, et al., Gynecologic Oncology, 50:89-93 (1993); Choy et al., Cancer, 71:3774-3778 (1993); Milas et al., Cancer Research, 54:3506-3510 (1994); and Joschko et al., Proceedings of the American Association for Cancer Research, 35:647 (1994). Although the reported data


suggests that taxol is an effective radiosensitizer, recent data from our laboratory leads us to question whether the reported data has been misinterpreted. In any event, a need continues to exist for compounds which
5 possess antitumor activity and which are more potent radiosensitizers and thus, can be administered at lower doses to reduce toxic side effects.

Summary of the Invention

Among the several objects of the invention,
10 therefore, may be noted the provision of a novel class of compounds and pharmaceutical preparations containing them which possess antitumor activity and which are potent radiosensitizing agents for cancer radiation therapy. Also among the objects of the invention are methods for
15 the use of such compounds and pharmaceutical preparations in warm-blooded animals in radiation therapy.

Briefly, therefore, the present invention is directed to taxanes comprising one or more electron-affinic moieties. Such compounds provide greatly
20 enhanced radiosensitization of tumors and reduced toxic side effects to normal body tissues at a given dosage as compared to conventional radiosensitization agents. The electron-affinic moiety may be attached directly, or indirectly through a linker to one of the ring atoms of
25 the taxane or to one of the C13 side chain atoms. For example, the electron-affinic moiety may be attached to the C2, C4, C7, C9, C10, C14, C3' or C5' carbon of a taxane corresponding to the structure:

4



wherein

M comprises ammonium or is a metal;

R₁ is hydrogen or hydroxy;

5 R₂ is -OT₂, -OCOZ₂, -OCOOZ₂, RSG₁ or RSG₂;

R₄ is -OT₄, -OCOZ₄, -OCOOZ₄, RSG₁ or RSG₂;


R₇ is hydrogen, halogen, -OT₇, -OCOZ₇, -OCOOZ₇,
RSG₁ or RSG₂;

10 R₉ is hydrogen, keto, -OT₉, -OCOZ₉, -OCOOZ₉, RSG₁
or RSG₂;

R₁₀ is hydrogen, keto, -OT₁₀, -OCOZ₁₀, -OCOOZ₁₀,
RSG₁ or RSG₂;

R₇, R₉, and R₁₀ independently have the alpha or
beta stereochemical configuration;

15 R₁₃ is hydroxy, protected hydroxy, keto, MO- or



R₁₄ is hydrogen, hydroxy, protected hydroxy,
RSG₁ or RSG₂;

20 T₂, T₄, T₇, T₉, and T₁₀ are independently hydrogen
or hydroxy protecting group;

X₁ is -OX₆;

X₂ is hydrogen, hydrocarbon, heterosubstituted
hydrocarbon, heteroaryl or heterosubstituted heteroaryl;

X_3 and X_4 are independently hydrogen, hydrocarbon, heterosubstituted hydrocarbon, heteroaryl, heterosubstituted heteroaryl or RSG_1 ;

X_5 is $-X_{10}$, $-OX_{10}$, $-SX_{10}$, or $-NX_8X_{10}$;

5 X_6 is hydrogen, hydrocarbon, heterosubstituted hydrocarbon, heteroaryl, heterosubstituted heteroaryl or hydroxy protecting group or a functional group which increases the water solubility of the taxane derivative;

10 X_8 is hydrogen, hydrocarbon, heterosubstituted hydrocarbon, RSG_1 or RSG_2 ;

X_{10} is hydrocarbon, heterosubstituted hydrocarbon, heteroaryl, heterosubstituted heteroaryl, RSG_1 or RSG_2 ;

15 Z_2 , Z_4 , Z_7 , Z_9 and Z_{10} are independently hydrocarbon, heterosubstituted hydrocarbon, heteroaryl or heterosubstituted heteroaryl;

RSG_1 is an electron-affinic moiety;

RSG_2 is $-L-(RSG_1)_n$;

20 L is a linker comprising a chain of 1 to 30 atoms in the chain, the atoms being selected from the group consisting of C, O, N, S, Si, and P; and

n is an integer greater than or equal to 1.

The invention is also directed to pharmaceutical compositions for radiosensitizing tumor cells which contain a radiosensitizing amount of the above described taxanes or a pharmaceutically acceptable salt thereof in admixture with a pharmaceutically acceptable carrier.

30 The present invention is further directed to a process for radiosensitizing tumor cells. The process comprises administering a radiosensitizing amount of the pharmaceutical composition described above to the tumor cells. Related thereto, a method is also provided for killing tumor cells in a warm-blooded animal which includes the steps of administering to the warm-blooded animal a pharmaceutical composition as described above in

an amount effective to radiosensitize the tumor cells, followed by, after a time interval sufficient to enhance radiosensitization of the tumor cells, irradiating the tumor cells with a dose of radiation effective to kill
5 the tumor cells.

Other objects and features will be in part apparent and in part pointed out hereinafter.

Brief Description of the Drawings

Fig. 1 is a graph depicting in vitro
10 chemotherapeutic activity of taxoltere metro, taxol and taxoltere pnip on CHO cells for the studies set forth in Example 4.1.

Fig. 2 is a graph depicting in vitro
chemotherapeutic activity of taxoltere, taxol and
15 taxoltere pnip on HCT 116 cells for the studies set forth in Example 4.1.

Fig. 3 is a graph depicting in vitro
chemotherapeutic radiosensitization of taxoltere metro
and taxoltere pnip on CHO cells for the studies set forth
20 in Example 4.2.

Fig. 4 is a graph depicting in vitro
chemotherapeutic radiosensitization of taxoltere metro
and taxoltere pnip on HCT 116 cells for the studies set
forth in Example 4.2.

25 Fig. 5 is a graph depicting in vivo dose-response curves for taxoltere metro and taxol for the studies set forth in Example 4.4.

Fig. 6 is a graph depicting in vivo
chemotherapeutic activity of taxoltere metro and
30 taxoltere pnip at 40% of LD₅₀ for the studies set forth in Example 4.4.

Fig. 7 is a graph depicting the
chemotherapeutic effect of low-dose multi-treatment (Q7D
x 4) with taxol and its analogs on MTG-B mammary tumors
35 for the studies set forth in Example 4.4.

Fig. 8 is a graph depicting the chemotherapeutic effects of low-dose multi-treatment (Q7D x 4) with taxol and its analogs on MTG-B mammary tumors for the studies set forth in Example 4.4.

5 Figs. 9 and 10 are graphs depicting the chemotherapeutic effects of low-dose multi-treatments (Q11D x 4) with taxol and its analogs on MTG-B mammary tumors for the studies set forth in Example 4.4.

10 Fig. 11 is a graph depicting in vivo chemotherapeutic radiosensitization of taxoltere metro, taxoltere pnip and taxol on MTG-B mammary tumors (i.p., single dose) for the studies set forth in Example 4.5.

15 Fig. 12 is a graph depicting the effects of ip taxol, taxoltere metro and taxoltere pnip +/-RT on MTG-B mammary tumors for the studies set forth in Example 4.5.

 Figs. 13 and 14 are graphs depicting the effects of taxoltere pnip on MTG-B mammary tumors (i.v., single dose, 24% LD50) for the studies set forth in Example 4.5.

20 Fig. 15 is a graph depicting the cure rate for taxoltere pnip (i.p., single dose) for the studies set forth in Example 4.6.

25 Figs. 16 and 17 are graphs depicting the cure rate for taxoltere pnip (i.v., single dose) for the studies set forth in Example 4.6.

 Fig. 18 is a graph depicting the cure rate for taxoltere pnip +/- RT on MTG-B mammary tumors in vivo as a function of drug dose for the studies set forth in Example 4.6.

30 Detailed Description of the Preferred Embodiments

 Surprisingly, it has been discovered that taxanes containing electron-affinic substituents exhibit significantly greater potency than taxol as a radiosensitizing agent. As a result, such increased
35 potency permits the administration of much lower dosages


of these compounds for the same or even greater radiosensitization of tumor cells, allowing for a concomitant reduction in toxic side effects on healthy tissue for any particular dosage level required to effectively radiosensitize the tumor cells.

The radiosensitizing groups or moieties described herein (RSG_1 and RSG_2) impart electron affinity to the compounds with which they are associated. This novel class of potent radiosensitizers comprise taxanes containing at least one, and optionally two or more electron-affinic moieties. In general, the radiosensitizing moieties contain electron-affinic groups which fall into one of four groups: (i) carbocyclic or heterocyclic aromatic moieties which possess one or more carbonyl, trifluoromethyl, halogen, nitro, sulfonyl, sulfinyl, phosphoryl, oxide or cyano groups, (ii) heterocyclic aromatic moieties containing two or more heteroatoms, (iii) metal complexes, and (iv) organo-metallic groups in which the metal is covalently bonded to carbon.

The carbocyclic or heterocyclic aromatic electron-affinic moieties contain one to three rings with a total of 5 to 15 ring atoms which are selected from the group consisting of C, N, S, O and P. Preferably, the carbocyclic or heterocyclic aromatic electron-affinic moieties contain one to two rings with one ring being presently most preferred. Representative carbocyclic aromatic electron-affinic moieties include phenyl and naphthyl groups containing one or more nitro, halogen, carbonyl or sulfonyl substituents, with nitro-substituted phenyl being a preferred carbocyclic aromatic electron-affinic moiety. Representative heterocyclic aromatic electron-affinic moieties include imidazoles, triazoles, pyridines, benzamides, nicotinamides, benzotriazine oxides, furans, thiophenes, oxazoles and thiozoles possessing one or more carbonyl, trifluoromethyl,

halogen, nitro, sulfonyl, sulfinyl, phosphoryl, oxide or cyano groups, and preferably at least one nitro group.

Nitroimidazole and nitrotriazole heterocyclic aromatic electron-affinic moieties which may be
 5 incorporated into the radiosensitizing agents of the present invention include 2-nitroimidazol-1-yl and 3-nitro-1,2,4-triazol-1-yl and other nitroimidazoles and nitrotriazoles which correspond to the following structures:




wherein E_1 is alkyl or fluoroalkyl. The preparation and use of radiosensitizing agents incorporating these and other nitroimidazoles and nitrotriazoles is described in Suzuki et al., U.S. Patent Nos. 4,945,102 and 5,064,849;
 15 Kagiza et al., U.S. Patent Nos. 4,927,941, 4,977,273 and 5,304,654; Suto, U.S. Patent No. 4,954,515 and 5,036,096; Suto et al., U.S. Patent No. 4,797,397; Papadopolou-Rosenzweig et al., U.S. Patent No. 5,294,715; Beylin et al., U.S. Patent No. 5,342,959.

20 Benzamide and nicotinamide heterocyclic aromatic electron-affinic moieties which may be incorporated into the radiosensitizing agents of the present invention include

- 5-hydroxynicotinamide;
- 25 5-nitronicotinamide;
- 5-(2,3-dihydroxypropoxy)nicotinamide;
- 5-aminonicotinamide;
- 5-(2-methoxyethylamino)nicotinamide;
- 5-acetamidonicotinamide;
- 30 3-hydroxy thiobenzamide;

3-[(2-hydroxyethoxy)acetamido]benzamide;
 3-(2,3 dihydroxy-n-propoxy)-4-methoxybenzamide;
 3-(2,3 dihydroxy-n-propoxy)-4-methylbenzamide;
 4-(2,3 dihydroxy-n-propoxy)-3-methoxybenzamide;

- 5 and other benzamides and nicotinamides which correspond to the following structures:




wherein X^1 is O or S; Y_1 is H, lower alkyl, lower alkoxy, acetoxy, or acetamido; Y_2 is -OR, -SR, -NHR, -NO₂,
 10 -O(CO)R, -NH(CO)R, -O(SO)R, or -O(POR)R; Y_3 is H, Z_1 , -OR, -SR, -NHR, -O(CO)R, -NH(CO)R, -O(SO)R, or -O(POR)R; and R is hydrogen or hydrocarbon which may be optionally substituted and interrupted by an ether (-O-) linkage.
 The preparation and use of radiosensitizing agents
 15 incorporating these and other benzamides and nicotinamides is described in Lee et al., U.S. Patent Nos. 5,032,617, 5,041,653 and 5,175,287.

Benzotriazine oxide heterocyclic aromatic electron-affinic moieties which may be incorporated into
 20 the radiosensitizing agents of the present invention include

3-hydroxy-1,2,4-benzotriazine-1,4-dioxide;
 3-amino-7-trifluoro-1,2,4-benzotriazine-1-oxide;
 25 3-amino-7-decyl-1,2,4-benzotriazine-1-oxide;
 3-amino-7-carbamyl-1,2,4-benzotriazine-1-oxide;
 7-acetyl-3-amino-1,2,4-benzotriazine-1-oxide;
 7-chloro-3-hydroxy-1,2,4-benzotriazine-1,4-dioxide;


7-nitro-3-amino-1,2,4-benzotriazine-1,4-dioxide; and other benzotriazine oxides corresponding to the structure:




5 wherein Y_4 is H, substituted or unsubstituted lower hydrocarbon, or alkanoyl; m is 0 or 1; and Y_5 and Y_6 are independently hydrogen, nitro, halogen, morpholino, pyrrolidino, piperidino, substituted or unsubstituted hydrocarbon, $-NH_2$, $-NHR'$, $-NR'R'O(CO)R'$, $-NH(CO)R'$,
 10 $-O(SO)R'$, or $-O(POR')R'$ in which R' is substituted or unsubstituted hydrocarbon. The preparation and use of radiosensitizing agents incorporating these and other benzotriazine oxides is described in Lee et al. U.S. Patent No. 5,175,287.

15 The metal complex electron-affinic moieties preferably comprise Pt^{2+} , Co^{3+} , Fe^{2+} , Fe^{3+} , Pd^{2+} , Cu^{2+} , Ti^{4+} , or Zr^{4+} as the metal and generally fall into two subgroups: (a) metal complexes of the carbocyclic and heterocyclic aromatic electron-affinic moieties discussed
 20 above, and (b) metal complexes of bidentate ligands comprising nitrogen, carbon or sulfur. In general, metal complexes of bidentate ligands correspond to the formula $-BM^kX_k$ wherein B is a bidentate ligand containing nitrogen, carbon or sulfur, M^k is a metal, X is an anionic ligand such as Cl^- or OAc^- , and k is 1-4. Exemplary
 25 bidentate ligands include:

12




Electron-affinic metal complexes which may be incorporated into the radiosensitizing agents of the present invention include compounds of the formula:



wherein n is 1 or 2, and wherein when n is 2, X^{M} is a monovalent biologically acceptable anion, and when j is 1, X^{M} is a divalent biologically acceptable anion; each R^{n} is independently H or alkyl, or both R^{n} 's together are a piperidino or morpholino residue; Q is a radiosensitizing ligand selected from a mononitro-substituted imidazole, a mononitro-substituted pyrazole, a mononitro-substituted thiazole and a mononitro-substituted isothiazole; and Y^- is a physiologically acceptable anion. These

15 heterocycles may optionally be substituted by an alkyl, amino substituted alkyl, hydroxy, alkoxy or amino group. In addition, if the heterocycle is pyrazole or imidazole, a ring nitrogen may be substituted by alkyl or alkoxy or hydroxy substituted alkyl and wherein one or two

20 methylenes of the alkyl may be replaced by oxygen. In a preferred embodiment, Q is one of the following:




wherein R^1 is alkyl optionally containing an amino substituent, $-\text{OR}^3$, or $-\text{N}(\text{R}^3)_2$, wherein R^3 is H or lower alkyl; R^2 is alkyl or 1-8 carbons substituted by one or more $-\text{OR}^3$ and wherein one or two methylenes may be

25

replaced by oxygen and each m is independently 0 or 1. The preparation and use of radiosensitizing agents incorporating these metal complexes is described in Skov et al. U.S. Patent Nos. 4,921,963 and 5,026,694.


5 Other electron-affinic metal complexes which may be incorporated into the radiosensitizing agents of the present invention may be made by reacting an organic or inorganic platinum compound such as an alkali metal tetrahaloplatinate or cis-bis(acetonitrile)dichloro-
10 platinum (II) with rhodamine 123 or other (+)-charged rhodamine or the like, for example, a cyanine dye such as 3,3'-diethylthiadibocyanine iodide or other (+)-charged cyanine dyes as described in U.S. Patent No. 5,196,413.

15 Other electron-affinic metal complexes which may be incorporated into the radiosensitizing agents of the present invention include include Cu(II) compounds selected from compounds having the formula:



20 wherein A^c represents a bidentate heteroaromatic ligand containing neutral nitrogen donor atoms; B^c represents a bidentate ligand containing neutral or negatively charged oxygen donor atoms; X^c and Y^c are the same or different neutral or negatively charged monodentate ligands; and Z¹
25 and Z² represent the charge on the complex. The preparation and use of radiosensitizing agents incorporating these metal complexes is described in Abrams et al. U.S. Patent No. 5,100,885.

30 Other electron-affinic metal complexes which may be incorporated into the radiosensitizing agents include Co(III) or Fe(III) compounds a formula corresponding to one of the following formulas:



35 wherein n has a value of 3 or 4; N is an uncharged nitrogen donor atom that is contained within a ligand; X^F represents an anionic ligand; and y represents the charge

on the complex; A^F represents a bidentate or tetradentate negative ligand containing N or O donor atoms; D^1 and D^2 represent the same or different monodentate ligands; q represents a positive or negative charge on the complex; Z^F represents a chelating mononegative negative ligand; T^1 and T^2 , which may be the same or different, represent mono-negative tridentate ligands. The preparation and use of radiosensitizing agents incorporating these metal complexes is described in U.S. Patent No. 4,727,068.







The organometallic electron-affinic moieties are aliphatic or aromatic mercury radicals. The preparation and use of radiosensitizing agents incorporating mercury containing entities is described in Shenoy et al., Cancer Investigation, 10(6):533-551 (1992) and Bruce et al., Radiation Res., 24:473-481 (1965).

The electron-affinic moieties may be directly attached to one of the carbons of the A, B, or C rings of the taxane or indirectly attached via a linker. The linker comprises a chain of 0 to 30 atoms in the chain, with approximately 10 or less being preferred. The chain atoms are selected from the group consisting of C, O, N, S, Si, and P and are preferably C, N or O. The linker may be linear or cyclic, branched or unbranched, and may contain as substituents, one or more P, C, O, N, S, H, Si or halogen-containing substituents. Exemplary linker substituents include silyls, ethers, thioethers, esters, thioesters, amides, thioamides, amines, alcohol, alkyl, aryl, carbonyl, sulfonyl, phosphoryl, and halogen substituents.





Preferably, the linker comprises a hydrocarbon segment consisting of 1 to 6 carbon atoms. It may additionally comprise a carbonyl, ester, thioester, amide, carbonate, thiocarbonate, carbamate, or ether segment. If a non-hydrocarbon segment is included; the non-hydrocarbon segment preferably comprises one or more

ether, carbonate or carbonyl moieties as the non-hydrocarbon segment.

For purposes of illustration, a series of radicals comprising linkers and electronic-affinic moieties falling within the scope of the present invention is set forth as follows:



16



5


wherein h is 1-3, R⁴ is H, hydrocarbon or substituted hydrocarbon, and R⁵ is hydrocarbon or substituted hydrocarbon. In other embodiments, the carbonyl or ester linkage of the above structures may be replaced by

10 thioester or amide linkages. In addition, many of these


radicals may serve as ligands for the previously identified metal species.

The radiosensitizing compounds of the present invention are prepared by linking the electron-affinic moiety to the C2, C4, C7, C9, C10, C14, C3', or C5' carbons of a taxane. The starting material may be 10-deacetyl baccatin III, baccatin III, or another naturally occurring taxane such as 14-hydroxy-10-deacetylbaccatin III. Alternatively, the taxane may be synthesized from commodity chemicals as set forth in PCT Patent Application No. WO 95/03265.

Taxanes having C13 side chains which incorporate electron-affinic moieties at C3' and/or C5' may be prepared through the use of β -lactams having the desired substituents and reacting the β -lactam and a C13 metal or ammonium alkoxide of a suitably substituted taxane as more fully described in U.S. Patent 5,430,160. The β -lactams have the following structural formula:



wherein X_1 is protected hydroxy, and $X_2 - X_5$ are as previously defined. Preferably, the alkoxide has the tetracyclic taxane nucleus and corresponds to the structural formula:




wherein M is a metal or tetraalkylammonium and R₁, R₂, R₄, R₇, R₉, R₁₀, and R₁₄ are as previously defined.

The electron-affinic moieties can be attached to the C2, C4, C7, C9, C10, and C14 positions of a taxane by a variety of methods. For purposes of illustration, the attachment method will first be described with respect to the C7 position. As will be described elsewhere herein, however, these same methods can be used for the other positions.





Metronidazole, a well known radiosensitizer, and other electron-affinic moieties can be attached via a carbonate linkage by treating baccatin III 2 with carbonyl diimidazole to produce the 7-carboxylimidazolide 2a, and reacting the product in situ with metronidazole at higher temperature to provide the 7-carboxylmetronidazolide 3. A C13 side chain can then be attached to 7-carboxylmetronidazolide 3 by treating it with lithium hexamethyldisilazide and β -lactam 4, and, after treatment with HF, to yield a taxane which we have named taxoltere metro 5. This reaction sequence is summarized in the following reaction scheme:

19




This method can be used for the preparation of a series of similar radiosensitizing taxanes having different radiosensitizing groups. The intermediate 7-carbonylimidazolidine **2a** reacts smoothly with alcohols to provide the desired radiosensitizing taxanes in which the radiosensitizing group is attached via a carbonate linkage, as in taxoltere metro. For example, alcohols **6**

through 9 will react with the carbonylimidazolidine substituent of 2a to yield four other taxanes having radiosensitizing groups linked to C7.




- 5 Attachment of the radiosensitizing group is then followed by attachment of a side chain at C13 in the same manner as it was accomplished for taxoltere metro.


The attachment of a metal atom or metal complex tethered at the C7 position of the taxane core can be accomplished by reacting an allylchloroformate with a taxane having an available C7 hydroxy group and a protected 2' hydroxy group to produce derivative 10 wherein X₃ and X₅ are as previously defined and P is a hydroxy protecting group. Hydroboration of the allyl carbonate substituent followed by treatment of the borane with mercuric acetate and sodium chloride and deprotection of the C2' hydroxy group gives the mercury derivative 11, a good radiosensitizer.




21





The ester analog of 11 can be prepared by the direct acylation of the C7 hydroxyl group of a 2' hydroxy protected taxane with an acid chloride to produce ester 12. Hydroboration of the allyl ester followed by treatment of the borane with mercuric acetate and sodium chloride gives the ester analog of mercury derivative 11.





Similar chemistry can be used to attach a bidentate ligand to the C7 position, and the metal complex (e.g., platinum) of the bidentate ligand can then be prepared by introducing an appropriate metallic reactant (e.g., PtCl_2 or $\text{PtCl}_2(\text{SMe}_2)_2$). The bidentate ligand can also incorporate an electron-affinic ligand, and the Pt (II) complex of 13 can be made in this way.




The length and nature of the linker between the taxane and the electron-affinic group may be altered. It is possible to prepare analogs in which the electron-affinic group is both closer and further away from the taxane than it is in taxoltere metro. An analog with a longer linker can easily be synthesized by incorporating a dicarboxylic acid diester instead of the carbonate between the taxane and metronidazole, e.g., 14. The *p*-nitrobenzyl ether 15 and the corresponding *p*-nitrobenzoate are radiosensitizing taxanes in which the electron-affinic group is very close to the taxane. Alternatively, hydroboration of 10 gives an alcohol, the mesylate of which reacts with, for example, 2-nitroimidazole to provide 16, and the epoxide derived by peracid treatment of 10 reacts with, for example, 2-nitroimidazole to provide 17. Ester analogs of 16 and 17 can be similarly prepared starting from 12.




23



It is also possible to prepare radiosensitizing taxanes having multiple radiosensitizing groups attached to a single linker. For example, reaction of 2-nitroimidazole with glycidyl chloride at somewhat elevated temperature provides alcohol 18, which then reacts with 2a and with β -lactam 4 to give radiosensitizing taxane 19.



10



Electron-affinic moieties can be attached to the C10 position of a taxane possessing a C10 hydroxy group, such as 10-DAB, by the methods discussed for attaching the electron-affinic moieties to the C7 position. Taxanes having other C10 substituents described herein may be prepared as more fully described in PCT Patent Application WO 94/15599 and other literature references. For example, taxanes having a C10 keto substituent can be prepared by oxidation of 10-desacetyl taxanes. Taxanes which are dihydro substituted at C10 can be prepared by reacting a C10 hydroxy or acyloxy substituted taxane with samarium diiodide. Taxanes having acyloxy substituents other than acetate can be prepared by reacting the C10 hydroxy substituent of 10-deacetyl baccatin III with any standard acylating agent such as anhydrides, acid chlorides, acyl imidazoles or other activated carboxyl derivatives. Taxanes having a C10 carbonate substituent can be prepared by using an analogous chloroformate instead of the acid chloride.


Electron-affinic moieties can be attached to the C9 position of a taxane possessing a C9 hydroxy group by the methods discussed for attaching the electron-affinic moieties to the C7 position. As more fully described in PCT Patent Application WO 94/20088, the C9
25 the C9 keto substituent of taxol, 10-DAB, baccatin III or can be selectively reduced to yield the corresponding C9

β -hydroxy derivative with a borohydride, preferably tetrabutylammonium borohydride (Bu_4NBH_4) or triacetoxy-borohydride. The C9 β -hydroxy derivative can then be protected at C7 with a hydroxy protecting group and the C9 hydroxy group can be acylated following the methods described herein for acylation of the C7 hydroxy group.

Alternatively, reaction of 7-protected-9 β -hydroxy derivative with KH causes the acetate group (or other acyloxy group) to migrate from C10 to C9 and the hydroxy group to migrate from C9 to C10, thereby yielding a 10-desacetyl derivative, which can be acylated as described elsewhere herein.

As more fully described in PCT Patent Application WO 94/17050, C7 dihydro and other C7 substituted taxanes can be prepared by tin hydride reduction of the C7 xanthate. C7 fluoro-substituted taxanes can be prepared by treatment of C13-triethylsilyl-protected baccatin III with 2-chloro-1,1,2-trifluorotriethylamine at room temperature in THF solution. Other baccatin derivatives with a free C7 hydroxyl group behave similarly. Alternatively, 7-chloro baccatin III can be prepared by treatment of baccatin III with methanesulfonyl chloride and triethylamine in methylene chloride solution containing an excess of triethylamine hydrochloride. Taxanes having C7 acyloxy substituents can be prepared as set forth in the following reaction scheme. 7,13-protected 10-oxo-derivative is converted to its corresponding C13 alkoxide by selectively removing the C13 protecting group and replacing it with a metal such as lithium. The alkoxide is then reacted with a β -lactam or other side chain precursor. Subsequent hydrolysis of the C7 protecting groups causes a migration of the C7 hydroxy substituent to C10, migration of the C10 oxo substituent to C9, and migration of the C9 acyloxy substituent to C7.

26




Taxanes having alternative C2 and/or C4 esters or carbonates which optionally may contain an electron-affinic moiety as described elsewhere herein can be prepared using baccatin III and 10-DAB as starting materials. The C2 and/or C4 esters of baccatin III and 10-DAB can be selectively reduced to the corresponding alcohol(s) using reducing agents such as LAH or Red-Al, and new esters can thereafter be substituted using standard acylating agents such as anhydrides and acid chlorides in combination with an amine such as pyridine, triethylamine, DMAP, or diisopropyl ethyl amine.

Alternatively, the C2 and/or C4 alcohols may be converted to new C2 and/or C4 esters through formation of the corresponding alkoxide by treatment of the alcohol with a suitable base such as LDA followed by an acylating agent such as an acid chloride. The corresponding carbonates can be prepared by substituting a chloroformate for the analogous acid chloride.


Baccatin III and 10-DAB analogs having different substituents at C2 and/or C4 can be prepared as set forth in Reaction Schemes C₂-1 to C₂-5. To simplify the description, 10-DAB is used as the starting material and only the ester products are shown. It should be understood, however, that other starting materials and reactants may be substituted to yield the other C2 and C4 substituted compounds disclosed herein.

In the Reaction Scheme C₂-1, protected 10-DAB is converted to the triol 21 with lithium aluminum hydride. Triol 21 is then converted to the corresponding C4 ester using Cl₂CO in pyridine followed by a nucleophilic agent (e.g., Grignard reagents or alkyllithium reagents) wherein Z₂ is as defined elsewhere herein.

28

Reaction Scheme C₂-1


- 5 Deprotonation of triol 21 with LDA followed by introduction of an acid chloride selectively gives the C4 ester. For example, when acetyl chloride was used, triol 21 was converted to 1,2 diol 24 as set forth in Reaction Scheme C₂-2 wherein Z₄ is as defined elsewhere herein.

Reaction Scheme C₂-2

Triol **21** can also readily be converted to the 1,2 carbonate **22**. Acetylation of carbonate **22** under vigorous standard conditions provides carbonate **25** as described in Reaction Scheme C₂-3; addition of


5 alkylolithiums or Grignard reagents to carbonate **22** provides the C2 ester **24** having a free hydroxyl group at C4 as set forth in Reaction Scheme C₂-1.

Reaction Scheme C₂-3




10 As set forth in Reaction Scheme C₂-4, other C4 substituents can be provided by reacting carbonate **22** with an acid chloride and a tertiary amine to yield carbonate **26** which is then reacted with alkylolithiums or Grignard reagents to provide 10-DAB derivatives **27** having


15 new substituents at C2 wherein Z₂ and Z₄ are as defined elsewhere herein.

Reaction Scheme C₂-4

Alternatively, baccatin III may be used as a starting material and reacted as shown in Reaction Scheme C₂-5. After being protected at C7 and C13, baccatin III is reduced with LAH to produce 1,2,4,10 tetraol 29. Tetraol 29 is converted to carbonate 30 using Cl_2CO and pyridine, and carbonate 30 is acylated at C10 with an acid chloride and pyridine to produce carbonate 31 (as shown) or with acetic anhydride and pyridine (not shown). Acetylation of carbonate 31 under vigorous standard conditions provides carbonate 32 which is then reacted with alkyl lithiums to provide the baccatin III derivatives 33 having new substituents at C2 and C10 wherein Z_2 and Z_{10} are as defined elsewhere herein.

Reaction Scheme C₂-5

32



Taxanes having radiosensitizing groups at the C14 position, i.e., the point of attachment of R₁₄ as depicted elsewhere herein, may be prepared using the same or similar methods as those described elsewhere herein with respect to attaching radiosensitizing groups to the C7 position of the taxane. The starting material for these compounds may be, for example, a hydroxylated taxane (14-hydroxy-10-deacetylbaccatin III) which has been discovered in an extract of yew needles (C&EN, p 36-37, April 12, 1993). Derivatives of this hydroxylated taxane having the various C2, C4, C7, C9, C10, C3' and C5' functional groups described above may also be prepared by using this hydroxylated taxane. In addition, the C14 hydroxy group together with the C1 hydroxy group of 10-DAB can be converted to a 1,2-carbonate as described in C&EN or it may be converted to a variety of

esters or other functional groups as otherwise described herein in connection with the C2, C4, C9 and C10 substituents.

The taxane radiosensitizers of the present invention can be combined with various excipient vehicles and/or adjuvants well known in this art which serve as pharmaceutically acceptable carriers to permit drug administration in the form of, e.g., injections, suspensions, emulsions, tablets, capsules, and ointments. These pharmaceutical compositions, containing a radiosensitizing amount of the described substituted diamine compounds, may be administered by any acceptable means which results in the radiosensitization of tumor cells. For warm-blooded animals, and in particular, for humans undergoing radiotherapy treatment, administration can be oral, parenteral, subcutaneous, intravenous, intramuscular and/or intraperitoneal. To destroy tumor cells, the pharmaceutical composition containing the radiosensitizing diamines are administered in an amount effective to radiosensitize the tumor cells (in the range of 1 to 100 mg/kg for humans). The specific dosage administered will be dependent upon such factors as the general health and physical condition of the patient as well as his age and weight, the stage of the patient's disease condition, and the existence of any concurrent treatments.

After administration of the radiosensitizing composition to the tumor cells and the passage of a time interval sufficient to enhance radiosensitization of the tumor cells, the tumor cells are irradiated with a dose of radiation effective to destroy the tumor cells. Generally, the patient will receive a total radiation dosage of about 60 to 76 Gy over seven to eight weeks, each individual radiation dose to be given within approximately 1 to 4 hours after administration of the radiosensitizer. Such sequences of radiosensitization

treatments and irradiation are repeated as needed to abate and, optimally, reduce or eliminate, the spread of the malignancy.

Definitions

5 The "hydrocarbon" moieties described herein are organic compounds or radicals consisting exclusively of the elements carbon and hydrogen. These moieties include alkyl, alkenyl, alkynyl, and aryl moieties. These moieties also include alkyl, alkenyl, alkynyl, and aryl
10 moieties substituted with other aliphatic or cyclic hydrocarbon groups, such as alkaryl, alkenaryl and alkynaryl. Preferably, these moieties comprise 1 to 20 carbon atoms.

 The alkyl groups described herein are
15 preferably lower alkyl containing from one to six carbon atoms in the principal chain and up to 20 carbon atoms. They may be straight or branched chain and include methyl, ethyl, propyl, isopropyl, butyl, hexyl and the like. They may be substituted with aliphatic or cyclic
20 hydrocarbon radicals or hetero-substituted with the various substituents defined herein.

 The alkenyl groups described herein are preferably lower alkenyl containing from two to six carbon atoms in the principal chain and up to 20 carbon
25 atoms. They may be straight or branched chain and include ethenyl, propenyl, isopropenyl, butenyl, isobutenyl, hexenyl, and the like. They may be substituted with aliphatic or cyclic hydrocarbon radicals or hetero-substituted with the various substituents
30 defined herein.

 The alkynyl groups described herein are preferably lower alkynyl containing from two to six carbon atoms in the principal chain and up to 20 carbon atoms. They may be straight or branched chain and
35 include ethynyl, propynyl, butynyl, isobutynyl, hexynyl,

and the like. They may be substituted with aliphatic or cyclic hydrocarbon radicals or hetero-substituted with the various substituents defined herein.

The aryl moieties described herein contain from
5 6 to 20 carbon atoms and include phenyl. They may be hydrocarbon or heterosubstituted with the various substituents defined herein. Phenyl is the more preferred aryl.


The heteroaryl moieties described are
10 heterocyclic compounds or radicals which are analogous to aromatic compounds or radicals and which contain a total of 5 to 20 atoms, usually 5 or 6 ring atoms, and at least one atom other than carbon, such as furyl, thienyl, pyridyl and the like. The heteroaryl moieties may be
15 substituted with hydrocarbon, heterosubstituted hydrocarbon or hetero-atom containing substituents with the hetero-atoms being selected from the group consisting of nitrogen, oxygen, silicon, phosphorous, boron, sulfur, and halogens. These substituents include lower alkoxy
20 such as methoxy, ethoxy, butoxy; halogen such as chloro or fluoro; ethers; acetals; ketals; esters; heteroaryl such as furyl or thienyl; alkanoxy; hydroxy; protected hydroxy; acyl; acyloxy; nitro; amino; and amido.

The heterosubstituted hydrocarbon moieties
25 described herein are hydrocarbon moieties which are substituted with at least one atom other than carbon, including moieties in which a carbon chain atom is substituted with a hetero atom such as nitrogen, oxygen, silicon, phosphorous, boron, sulfur, or a halogen atom.
30 These substituents include lower alkoxy such as methoxy, ethoxy, butoxy; halogen such as chloro or fluoro; ethers; acetals; ketals; esters; heteroaryl such as furyl or thienyl; alkanoxy; hydroxy; protected hydroxy; acyl; acyloxy; nitro; amino; and amido.

The acyl moieties described herein contain hydrocarbon, substituted hydrocarbon or heteroaryl moieties.

The alkoxycarbonyloxy moieties described herein
5 comprise lower hydrocarbon or substituted hydrocarbon moieties.

The term "taxane" as used herein, denotes compounds containing the A, B and C rings (with numbering of the ring positions shown herein):



10

As used herein "Ac" means acetyl; "AIBN" means azo-(bis)-isobutyronitrile; "Ar" means aryl; "BMDA" means BrMgNiPr_2 ; "BOC" means butyloxycarbonyl; "BOM" means benzyloxymethyl; "10-DAB" means 10-desacetylbaaccatin III;
15 "DBU" means diazabicycloundecane; "DMAP" means p-dimethylamino pyridine; "DDQ" means dicyano-dichloroquinone; "DMF" means dimethylformamide; "Et" means ethyl; "FAR" means 2-chloro-1,1,2-trifluoro-triethylamine; "iPr" means isopropyl; "LAH" means lithium
20 aluminum hydride; "LDA" means lithium diisopropylamide; "LHMDS" means lithium hexamethyldisilazide; "LTMP" means lithium tetramethylpiperidide; "mCPBA" means metachloro-perbenzoic acid; "Me" means methyl; "MOP" means 2-methoxy-2-propyl; "Ms" means CH_3SO_2 -; "Ph" means phenyl;
25 "protected hydroxy" means -OP or -OT wherein P or T is a hydroxy protecting group; "py" means pyridine; "R" means lower alkyl unless otherwise defined; "Red-Al" means sodium bis(2-methoxyethoxy) aluminum hydride; "Swern" means $(\text{COCl})_2$, Et_3N ; "TASF" means tris(diethylamino)-
30 sulfoniumdifluorotrimethyl-silicate; "TBAF" means tetrabutylammonium fluoride; "tBu" and "t-Bu" means tert-

butyl; "TBS" means $\text{Me}_2\text{t-BuSi-}$; "TES" means triethylsilyl; "Tf" means $-\text{SO}_2\text{CF}_3$; "TMS" means trimethyl-silyl; "TPAP" means tetrapropylammonium perruthenate; and "Ts" means toluenesulfonyl. "Hydroxy protecting group" includes, but is not limited to, acetals having two to ten carbons, ketals having two to ten carbons, ethers such as methyl, t-butyl, benzyl, p-methoxybenzyl, p-nitrobenzyl, allyl, trityl, methoxymethyl, methoxyethoxymethyl, ethoxyethyl, tetrahydropyranyl, tetrahydrothiopyranyl, and trialkylsilyl ethers such as trimethylsilyl ether, triethylsilyl ether, dimethylarylsilyl ether, triisopropylsilyl ether and t-butyl dimethylsilyl ether; esters such as benzoyl, acetyl, phenylacetyl, formyl, mono-, di-, and trihaloacetyl such as chloroacetyl, dichloroacetyl, trichloroacetyl, trifluoroacetyl; and carbonates including but not limited to alkyl carbonates having from one to six carbon atoms such as methyl, ethyl, n-propyl, isopropyl, n-butyl, t-butyl; isobutyl, and n-pentyl; alkyl carbonates having from one to six carbon atoms and substituted with one or more halogen atoms such as 2,2,2-trichloroethoxymethyl and 2,2,2-trichloroethyl; alkenyl carbonates having from two to six carbon atoms such as vinyl and allyl; cycloalkyl carbonates having from three to six carbon atoms such as cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl; and phenyl or benzyl carbonates optionally substituted on the ring with one or more C_{1-6} alkoxy, or nitro. Other hydroxyl protecting groups may be found in "Protective Groups in Organic Synthesis" by T. W. Greene, John Wiley and Sons, 1981.

To further illustrate and explain the invention, several examples are presented below.


EXAMPLE 1

Preparation of 7-(metronidazoleoxycarbonyl)baccatin III
hn-4-99-4d-2-1

- To a solution of baccatin III (100 mg, 0.170 mmol) in anhydrous 1,2-dichloroethane (1 mL) under nitrogen was added 1,1'-carbonyldiimidazole (55 mg, 0.40 mmol) and the reaction mixture was warmed up to 60 °C and stirred at that temperature for 10 h at which time the reaction was complete. Metronidazole (292 mg, 1.72 mmol) was added (neat) and the mixture was refluxed under nitrogen atmosphere. The progress of the reaction was monitored by NMR. When the reaction was complete (approximately three days) the mixture was diluted with ethyl acetate and washed with saturated sodium bicarbonate, brine. The organic layer was separated, dried and concentrated. The crude mixture was purified by flash chromatography to give 81 mg (70%) of the 7-(metronidazole-oxycarbonyl)baccatin III: mp. 220-223 °C; $[\alpha]_{\text{Hg}}^{25} = -44.9$
- ¹H NMR (CDCl₃, 500 MHz) δ 8.10 (d, J=7.1 Hz, 2H, benzoate), 7.96 (s, 1H, imidazole), 7.62-7.46 (m, 3H, benzoate), 6.28 (s, 1H, H10), 5.61 (d, J=6.8 Hz, 1H, H2β), 5.49 (m, 1H, H7) 4.96 (d, J=8.2 Hz, 1H, H5), 4.85 (br s, 1H, H¹³) 4.62-4.43 (m, 4H, methylenes), 4.31 (d, J=8.5 Hz, 1H, H20α), 4.13 (d, J=7.5 Hz, 1H, H20β), 3.99 (d, J=6.8 Hz, 1H, H3), 2.59 (m, 1H, H6α), 2.50 (s, 3H, CH3-imidazole), 2.29 (m, 5H, 4Ac, H14's), 2.13 (s, 3H, 10Ac), 2.10 (br s, 3H, Me18) 2.06 (d, J=5 Hz, 1H, 13OH), 1.9 (m, 1H, H6β), 1.77 (s, 3H, Me19), 1.59 (s, 1H, 10H), 1.17 (s, 3H, Me17), 1.08 (s, 3H, Me16).

EXAMPLE 2

Preparation of N-debenzoyl-N-(t-butylcarbamoyl)-7-(metronidazoleoxycarbonyl) taxol. (Taxoltere metro)



5 hn-4-119-2

To a solution of 7-(metronidazoleoxycarbonyl) baccatin III (31 mg, 0.039 mmol) in 0.3 mL of THF at -45 °C was added dropwise 0.048 mL of a 1.00 M solution of lithium bis(trimethylsilyl)amide in THF. After 0.5 h at
 10 -45 °C, a solution of *cis*-1-(t-butoxycarbonyl)-3-triethylsilyloxy-4-phenylazetidin-2-one (75 mg, 0.20 mmol) in 0.3 mL of THF was added dropwise to the mixture. The solution was warmed to 0 °C and kept at that temperature for 1 h before 0.2 mL of a 10% solution of
 15 AcOH in THF was added. The mixture was partitioned between saturated aqueous NaHCO₃ and 60/40 ethyl acetate/hexane. Evaporation of the organic layer gave a residue which was purified by filtration through silica gel to give 45.0 mg of a mixture containing (2'R,3'S)-2'-
 20 triethylsilyl-N-debenzoyl-N-(t-butylcarbamoyl)-7-(metronidazoleoxycarbonyl) taxol and a small amount of the (2'S,3'R) isomer.


To a solution of 45.0 mg of the mixture obtained from the previous reaction in 1.5 mL of
 25 acetonitrile and 0.6 mL of pyridine at 0 °C was added 0.2 mL of 48% aqueous HF. The mixture was stirred at 0 °C for 3 h, then at 25 °C for 13 h, and partitioned between saturated aqueous sodium bicarbonate and ethyl acetate. Evaporation of the ethyl acetate solution gave 40.8 mg of

material which was purified by plug filtration and recrystallization from methanol/water to give 32.3 mg (79%) of N-debenzoyl-N-(t-butylcarbamoyl)-7-(metronidazoleoxy-carbonyl) taxol. m.p. 169-172 °C; $[\alpha]^{25}_{Na}$ -52 °C (0.0035, CHCl₃).

¹H NMR (CDCl₃, 300 MHz) δ 8.10 (d, J=7.1 Hz, 2H, benzoate), 7.96 (s, 1H, imidazole), 7.62-7.26 (m, 8H, benzoate, 3'-phenyl), 6.25 (s, 1H, H10), 6.18 (dd, J = 8.8, 8.8 Hz, 1H, H13), 5.65 (d, J=7.1 Hz, 1H, H2β), 5.44-5.21 (m, 3H, NH, H3', H2'), 4.91 (d, J=9.9 Hz, 1H, H5), 4.62 (m, 4H, methylenes), 4.44 (m, 1H, H7), 4.31 (d, J = 8.2 Hz, 1H, H20α), 4.15 (d, J = 8.2 Hz, 1H, H20β), 3.90 (d, J=7.1 Hz, 1H, H3), 3.35 (d, J=5.5 Hz, 1H, 2'-OH), 2.59 (m, 1H, H6α), 2.49 (s, 3H, CH₃-imidazole), 2.36 (s, 3H, 4Ac), 2.31 (m, 2H, H14), 2.13 (s, 3H, 10Ac), 1.95 (m, 1H, H6β), 1.88 (br s, 3H, Me18), 1.77 (s, 3H, Me19), 1.70 (s, 1H, 1OH), 1.34 (s, 9H, t-butyl), 1.23 (s, 3H, Me17), 1.15 (s, 3H, Me16).

EXAMPLE 3

Preparation of N-debenzoyl-N-(t-butylcarbamoyl)-3'-desphenyl-3'-(4-nitrophenyl) taxol. (Taxoltere p-nip)



hn-2-66-3

To a solution of 7-triethylsilyl baccatin III (120 mg, 0.171 mmol) in 1.2 mL of THF at -45 °C was added

dropwise 0.104 mL of a 1.63 M solution of nBuLi in hexane. After 0.5 h at -45 °C, a solution of *cis*-1-(*t*-butoxycarbonyl)-3-triethylsilyloxy-4-(4-nitrophenyl)azetidin-2-one (361 mg, 0.885 mmol) in 1.2 mL of THF was added dropwise to the mixture. The solution was warmed to 0 °C and kept at that temperature for 1 h before 1 mL of a 10% solution of AcOH in THF was added. The mixture was partitioned between saturated aqueous NaHCO₃ and 60/40 ethyl acetate/hexane. Evaporation of the organic layer gave a residue which was purified by filtration through silica gel to give 192 mg of a mixture containing (2'R,3'S)-2',7-(bis)triethylsilyl-N-debenzoyl-N-(*t*-butylcarbamoyl)-3'-desphenyl-3'-(4-nitrophenyl) taxol and a very small amount of the (2'S,3'R) isomer.

To a solution of 192 mg of the mixture obtained from the previous reaction in 11 mL of acetonitrile and 0.55 mL of pyridine at 0 °C was added 1.7 mL of 48% aqueous HF. The mixture was stirred at 0 °C for 3 h, then at 25 °C for 13 h, and partitioned between saturated aqueous sodium bicarbonate and ethyl acetate. Evaporation of the ethyl acetate solution gave 153 mg of material which was purified by flash chromatography to give 140 mg (91%) of N-debenzoyl-N-(*t*-butylcarbamoyl)-3'-desphenyl-3'-(4-nitrophenyl) taxol, which was recrystallized from methanol/water.

m.p. 172-173 °C; $[\alpha]_{\text{Na}}^{25} -54$ °C (c 0.0046, CHCl₃).

EXAMPLE 4

Biological Studies of Taxoltere Metro and Taxoltere p-nip

4.1. In vitro chemotherapeutic activity.

Chinese hamster ovary (CHO) and human colon carcinoma (HCT-116) cells were treated with different concentrations of taxol, taxoltere metro, or taxoltere p-nip at 37 °C for 24 hours. Cell survival was evaluated by the colony forming assay. As shown in Figures 1 and

2, both taxoltere metro and taxoltere p-nip are much more efficient than taxol in killing CHO and HCT-116 cells. At the 50% survival level, taxoltere metro is about 15 times, and taxoltere p-nip is about 45 times, more effective than taxol in killing CHO cells; measured at the 1% survival level, taxoltere metro is about 10 times, and taxoltere p-nip is about 30 times, more effective than taxol in killing HCT-116 cells. As illustrated in Figures 1 and 2, both taxoltere metro and taxoltere p-nip exhibit significantly stronger ability than taxol to kill both types of cells at every drug dose point.

4.2. In vitro chemotherapeutic radiosensitization. These studies were carried out as above except that cells were irradiated (General Electric Maxitron 300 at 250 kvp and 20 mA (HVL 20 mm Al filter; dose rate of 2 Gy/min) after two hours incubation. Figures 3 and 4 show the results of experiments in which cells were subjected to different radiation doses in the presence or absence of drugs. Both taxoltere metro and taxoltere p-nip strongly radiosensitize both CHO and HCT-116 cells, although taxol does not. For CHO cells, the sensitizer enhancement ratio (SER) is 2.3 for 100 nM taxoltere metro and 1.6 for taxoltere p-nip. For HCT-116 cells, taxoltere metro shows a SER of 1.2 at the (low) dose of 10 nM, and taxoltere p-nip has a SER of 1.5. At each radiation dose point, there is a significantly enhanced decrease in the surviving fraction for the taxoltere metro and taxoltere p-nip treated groups, but not for the taxol treated groups. HCT-116 cells are more sensitive to both taxoltere metro and taxoltere p-nip than CHO cells, hence lower concentrations of the drugs are required for a significant enhancement of radiation induced cell killing.

4.3. **In vivo drug toxicity.** Acute toxicity experiments were conducted on C3H/HeJ mice. The LD₅₀ (lethal dose to 50% of animals) values were determined by standard procedures described by Chan and Hayes (Chan, P.K. and Hayes, A.W. Principles and Methods for acute toxicity and eye irritancy. In Hayes, A.W., ed. Principles and Methods of Toxicity. 2nd Ed., New York, New York, Raven Press; 1989: 169220). The LD_{50/5} for i.p. taxoltere metro is 249.67 mg/kg, compared with 140.97 mg/kg for i.p. taxol. At high drug dose levels, the death of mice in the taxol treated groups occurred sooner than the death of mice in the taxoltere metro treated groups. Severe toxic symptoms such as the constriction of pupils and the contraction of erectile tissue of hair follicles (resulting in rough hair) were observed 24 hours after drug administration in the taxol treated groups, but not in the taxoltere metro treated groups. Obviously, the acute toxicity of taxoltere metro is significantly lower than that of taxol. The LD_{50/5} values are 79.13 mg/kg for i.p. administration of taxoltere p-nip and 134.16 mg/kg for i.v. administration of taxoltere p-nip. The data strongly demonstrate that taxoltere p-nip is significantly less toxic when administered i.v. than it is when administered i.p.

4.4. **In vivo chemotherapeutic activity.** Taxol, taxoltere metro, and taxoltere p-nip were administered at equitoxic doses using C3H/HeJ mice bearing a mouse mammary adenocarcinoma (MTG-B) in the right flank. As illustrated in Figure 5, taxol-metro is much more effective than taxol in increasing the life span of mammary tumor-bearing mice. The survival time for control mice was 13 days. At 40% of the LD_{50/5} dose, taxol increased the survival time by 31% to 17 days, and taxoltere metro increased the survival time by 123% to 29 days. Taxoltere p-nip (Figure 6) is substantially more

potent, increasing the life span by 1015% to 145 days (dose-response data not shown because only a single dose (40% of $LD_{50/5}$) was studied). Thus, under equitoxic conditions, taxoltere metro was 4 times, and taxoltere p-nip was 33 times, more effective than taxol.

Two experiments in which fractionated lower doses (20% of $LD_{50/5}$) of taxol, taxoltere metro and taxoltere p-nip were administered to MTG-B tumor bearing C3H/Hej mice have been carried out. In the first of these, drugs were administered on a q7d x 4 schedule (Figure 7). As shown in Figure 8, tumor doubling times are greatly extended by taxoltere metro and are even further extended by taxoltere p-nip on this administration schedule. In the second experiment (Figure 9), drugs were administered on a q11d x4 schedule. Using this schedule, tumor doubling times (shown in Figure 10) are again greatly extended, with taxoltere metro producing the better results. Efficacy is positively correlated with frequency of injection (i.e., smaller intervals between injections). On both schedules, taxoltere metro and taxoltere p-nip exhibit significantly stronger antitumor effects than taxol.

4.5. In vivo chemotherapeutic radiosensitization. MTG-B tumor bearing C3H/Hej mice received single equitoxic drug doses i.p. (40% of $LD_{50/5}$). One day later the mice were placed in a lead holder (with the tumor-bearing hind leg exposed), and the tumor was subjected to a 22 Gy Xray dose (Figure 11). As shown in Figure 12, tumor doubling times (TDT) of 3.5 days (control), 8.5 days (22 Gy Xray exposure alone), 14.5 days (taxol alone), 19.5 days (taxol plus 22 Gy Xray), 20.5 days (taxoltere metro alone), and 29.5 days (taxoltere metro plus 22 Gy Xray) were observed. Equally large effects were observed when mice were given taxoltere metro two hours prior to irradiation (data not

shown). Taxoltere p-nip alone (TDT 3D 36.5 days) is even more effective in delaying tumor growth (Figure 12), and the combination of taxoltere p-nip plus 22 Gy produced a TDT of 65 days.

5 Administration of taxoltere p-nip i.v. (24% of $LD_{50/5}$) (Figure 13), gave a tumor doubling time of 52 days (Figure 14), approximately equivalent to the results obtained from i.p. administration.

4.6. **Cure Rates.** The term "cure" is defined
10 by the U.S. National Cancer Institute as tumor-free survival for at least twice the survival time of control tumor-bearing mice, therefore we have used 28 days tumor-free to define a "cure" in this system. The cure rate is 40% for mice treated i.p. with taxoltere p-nip
15 alone at a single dose, and 75% for the combination of taxoltere p-nip and 22 Gy (Figure 15). Similarly, a single i.v. injection of taxoltere p-nip alone induces a cure rate of 17%, and an 83% cure rate was observed for the combination of taxoltere p-nip and 22 Gy (Figure 16),
20 even though this experiment was conducted with a lower dose (24% of $LD_{50/5}$). Although in this latter group one tumor recurred after 40 days, some treated mice have remained tumor-free for more than one year, and still survive.

25 As shown in Figure 17, administration of taxoltere p-nip iv at 40% of $LD_{50/5}$ produced a cure rate of 50%, and iv administration of taxoltere p-nip at 40% of $LD_{50/5}$ in combination with 22 Gy radiation produced a cure rate of 100%. Cure rate as a function of drug dose is
30 shown in Figure 18.

In view of the above, it will be seen that the several objects of the invention are achieved and other advantageous results attained.

35 As various changes could be made in the above compounds and methods without departing from the scope of

the invention it is intended that all matter contained in the above description shall be interpreted as illustrative and not in a limiting sense.


WHAT IS CLAIMED IS:

1. A compound comprising a taxane containing at least 2 electron-affinic radiosensitizing functional groups.

2. The compound of claim 1 wherein at least one of the radiosensitizing groups is a nitro-substituted carbocyclic or heterocyclic aromatic moiety which is attached to the C2, C4, C7, C9, C10 or C14 position of the taxane.

3. The compound of claim 1 wherein said radiosensitizing groups are independently selected from nitro-substituted carbocyclic and heterocyclic aromatic moieties and wherein at least one of said radiosensitizing groups is attached to the C2, C4, C7, C9, C10 or C14 position of the taxane.

4. A compound corresponding to the structure:




wherein

- M comprises ammonium or is a metal;
- R₁ is hydrogen or hydroxy;
- R₂ is -OT₂, -OCOT₂, -OCOT₂, RSG₁ or RSG₂;
- R₄ is -OT₄, -OCOT₄, -OCOT₄, RSG₁ or RSG₂;
- R₇ is hydrogen, halogen, -OT₇, -OCOT₇, -OCOT₇, RSG₁ or RSG₂;
- R₉ is hydrogen, keto, -OT₉, -OCOT₉, -OCOT₉, RSG₁ or RSG₂;

R_{10} is hydrogen, keto, $-OT_{10}$, $-OCOZ_{10}$, $-OCOOZ_{10}$,
RSG₁ or RSG₂;

R_7 , R_9 , and R_{10} independently have the alpha or
15 beta stereochemical configuration;

R_{13} is hydroxy, protected hydroxy, keto, MO- or



R_{14} is hydrogen, hydroxy, protected hydroxy,
RSG₁ or RSG₂;

20 T_2 , T_4 , T_7 , T_9 and T_{10} are independently hydrogen
or hydroxy protecting group;

X_1 is $-OX_6$;

X_2 is hydrogen, hydrocarbon, heterosubstituted
hydrocarbon, heteroaryl, or heterosubstituted heteroaryl;

25 X_3 is alkyl, heterosubstituted alkyl, alkenyl,
heterosubstituted alkenyl, alkynyl, heterosubstituted
alkynyl, phenyl, heteroaryl, or heterosubstituted
heteroaryl;

X_4 is hydrogen, alkyl, heterosubstituted alkyl,
30 alkenyl, heterosubstituted alkenyl, alkynyl,
heterosubstituted alkynyl, phenyl, heteroaryl, or
heterosubstituted heteroaryl;

X_5 is $-X_{10}$, $-OX_{10}$, $-SX_{11}$, or $-NX_8X_{11}$;

X_6 is hydrogen, hydrocarbon, heterosubstituted
35 hydrocarbon, heteroaryl, heterosubstituted heteroaryl,
hydroxy protecting group or a functional group which
increases the water solubility of the taxane derivative;

X_8 is hydrogen, hydrocarbon, heterosubstituted
hydrocarbon, RSG₁ or RSG₂;

40 X_{10} is alkyl, heterosubstituted alkyl, alkenyl,
heterosubstituted alkenyl, alkynyl, heterosubstituted

alkynyl, phenyl, heteroaryl, or heterosubstituted heteroaryl;

45 X_{11} is hydrocarbon, heterosubstituted hydrocarbon, heteroaryl, heterosubstituted heteroaryl, RSG₁ or RSG₂;

Z_2 , Z_4 , Z_7 , Z_9 and Z_{10} are independently hydrocarbon, heterosubstituted hydrocarbon, heteroaryl, or heterosubstituted heteroaryl;

50 RSG₁ is an electron-affinic moiety;

RSG₂ is -L-(RSG₁)_n;

L is a linker comprising a chain of 1 to 30 atoms in the chain, the atoms being selected from the group consisting of C, O, N, S, Si, and P; and

55 n is an integer greater than or equal to 1; provided that the compound contains at least one radiosensitizing group attached to the C2, C4, C7, C9, C10, C14, C3' or C5' position of the compound.

5. The compound of claim 4 wherein RSG₁ is an electron-affinic group selected from the group consisting of (i) carbocyclic and heterocyclic aromatic moieties which possess one or more carbonyl, trifluoromethyl, halogen, nitro, sulfonyl, sulfinyl, phosphoryl, oxide or cyano groups, (ii) heterocyclic aromatic moieties containing two or more heteroatoms, (iii) metal complexes, and (iv) organo-metallic groups in which the metal is covalently bonded to carbon.

6. The compound of claim 4 wherein RSG₁ is selected from the group consisting of imidazoles, triazoles, pyridines, benzamides, nicotinamides, benzotriazine oxides, furans, thiophenes, oxazoles and thiozoles possessing one or more carbonyl, trifluoromethyl, halogen, nitro, sulfonyl, sulfinyl, phosphoryl, oxide or cyano groups.

50

7. The compound of claim 4 wherein

R_1 is hydrogen or hydroxy;

R_2 is $-\text{OCOZ}_2$, RSG_1 , or RSG_2 ;


R_4 is $-\text{OCOZ}_4$, RSG_1 or RSG_2 ;

5 R_7 is hydrogen, halogen, $-\text{OT}_7$, $-\text{OCOZ}_7$, RSG_1 or RSG_2 ;

R_9 is hydrogen, keto, $-\text{OT}_9$, $-\text{OCOZ}_9$, RSG_1 or RSG_2 ;

R_{10} is hydrogen, keto, $-\text{OT}_{10}$, $-\text{OCOZ}_{10}$, RSG_1 or RSG_2 ;

10 R_{13} is



R_{14} is hydrogen, hydroxy or protected hydroxy;

T_2 , T_4 , T_7 , T_9 and T_{10} are independently hydrogen or hydroxy protecting group;

15 X_1 is $-\text{OX}_6$;

X_2 is hydrogen;

X_3 is alkyl, alkenyl, phenyl, heteroaryl, or heterosubstituted heteroaryl;

20 X_4 is hydrogen, hydrocarbon, heteroaryl, or heterosubstituted heteroaryl;

X_5 is $-\text{X}_{10}$, $-\text{OX}_{10}$, $-\text{SX}_{11}$, or $-\text{NX}_8\text{X}_{11}$;

X_6 is hydrogen or hydroxy protecting group;

X_8 is hydrogen, hydrocarbon, heterosubstituted hydrocarbon, RSG_1 or RSG_2 ;

25 X_{10} is alkyl, alkenyl, phenyl, heteroaryl, or heterosubstituted heteroaryl;

Z_2 , Z_4 , Z_7 , Z_9 and Z_{10} are independently hydrocarbon, heterosubstituted hydrocarbon, heteroaryl, or heterosubstituted heteroaryl;

30 RSG_1 is an electron-affinic moiety;

RSG_2 is $-\text{L}-(\text{RSG}_1)_n$;

L is a linker comprising a chain of 1 to 10 atoms in the chain, the atoms being selected from the group consisting of C, O, N, S, Si, and P; and

35


n is 1 or 2.

8. The compound of claim 4 wherein RSG₁ is a heterocyclic aromatic moiety containing two or more heteroatoms.

9. The compound of claim 4 wherein RSG₁ is a metal complex.

10. The compound of claim 4 wherein RSG₁ is selected from the group consisting imidazoles, triazoles, pyridines, benzamides, furans, thiophenes, oxazoles and thiozoles possessing one or more nitro groups.

11. A compound corresponding to the structure:



wherein

M comprises ammonium or is a metal;

5

R₁ is hydrogen or hydroxy;

R₂ is RSG₁ or RSG₂;

R₄ is -OT₄, -OCOOZ₄, -OCOOZ₄, RSG₁ or RSG₂;

R₇ is hydrogen, halogen, -OT₇, -OCOOZ₇, -OCOOZ₇, RSG₁ or RSG₂;


10

R₉ is hydrogen, keto, -OT₉, -OCOOZ₉, -OCOOZ₉, RSG₁ or RSG₂;

R₁₀ is hydrogen, keto, -OT₁₀, -OCOOZ₁₀, -OCOOZ₁₀, RSG₁ or RSG₂;

R_7 , R_9 , and R_{10} independently have the alpha or
 15 beta stereochemical configuration;

R_{13} is hydroxy, protected hydroxy, keto, MO- or



R_{14} is hydrogen, hydroxy, protected hydroxy,
 RSG₁ or RSG₂;

20 T_2 , T_4 , T_7 , T_9 and T_{10} are independently hydrogen
 or hydroxy protecting group;

X_1 is -OX₆;

X_2 is hydrogen, hydrocarbon, heterosubstituted
 hydrocarbon, heteroaryl, or heterosubstituted heteroaryl;

25 X_3 and X_4 are independently hydrogen,
 hydrocarbon, heterosubstituted hydrocarbon, heteroaryl,
 heterosubstituted heteroaryl or RSG₁;

X_5 is -X₁₀, -OX₁₀, -SX₁₀, or -NX₈X₁₀;

30 X_6 is hydrogen, hydrocarbon, heterosubstituted
 hydrocarbon, heteroaryl, heterosubstituted heteroaryl,
 hydroxy protecting group or a functional group which
 increases the water solubility of the taxane derivative;

X_8 is hydrogen, hydrocarbon, heterosubstituted
 hydrocarbon, RSG₁ or RSG₂;

35 X_{10} is hydrocarbon, heterosubstituted
 hydrocarbon, heteroaryl, heterosubstituted heteroaryl,
 RSG₁ or RSG₂;


40 Z_4 , Z_7 , Z_9 and Z_{10} are independently hydrocarbon,
 heterosubstituted hydrocarbon, heteroaryl, or
 heterosubstituted heteroaryl;

RSG₁ is an electron-affinic moiety;

RSG₂ is -L-(RSG₁)_n;


L is a linker comprising a chain of 1 to 30 atoms in the chain, the atoms being selected from the group consisting of C, O, N, S, Si, and P; and
 45 n is an integer greater than or equal to 1.

12. A compound corresponding to the structure:



wherein

- M comprises ammonium or is a metal;
 5 R₁ is hydrogen or hydroxy;
 R₂ is -OT₂, -OC(O)Z₂, -OC(O)OZ₂, RSG₁ or RSG₂;
 R₄ is RSG₁ or RSG₂;
 R₇ is hydrogen, halogen, -OT₇, -OC(O)Z₇, -OC(O)OZ₇,
 RSG₁ or RSG₂;
 10 R₉ is hydrogen, keto, -OT₉, -OC(O)Z₉, -OC(O)OZ₉, RSG₁
 or RSG₂;
 R₁₀ is hydrogen, keto, -OT₁₀, -OC(O)Z₁₀, -OC(O)OZ₁₀,
 RSG₁ or RSG₂;
 R₇, R₉, and R₁₀ independently have the alpha or
 15 beta stereochemical configuration;
 R₁₃ is hydroxy, protected hydroxy, keto, MO- or



R₁₄ is hydrogen, hydroxy, protected hydroxy,
 RSG₁ or RSG₂;

20 T_2 , T_4 , T_7 , T_9 and T_{10} are independently hydrogen or hydroxy protecting group;

X_1 is $-OX_6$;

X_2 is hydrogen, hydrocarbon, heterosubstituted hydrocarbon, heteroaryl, or heterosubstituted heteroaryl;

25 X_3 and X_4 are independently hydrogen, hydrocarbon, heterosubstituted hydrocarbon, heteroaryl, heterosubstituted heteroaryl, or RSG_1 ;

X_5 is $-X_{10}$, $-OX_{10}$, $-SX_{10}$, or $-NX_8X_{10}$;

30 X_6 is hydrogen, hydrocarbon, heterosubstituted hydrocarbon, heteroaryl, heterosubstituted heteroaryl, hydroxy protecting group or a functional group which increases the water solubility of the taxane derivative;

X_8 is hydrogen, hydrocarbon, heterosubstituted hydrocarbon, RSG_1 or RSG_2 ;

35 X_{10} is hydrocarbon, heterosubstituted hydrocarbon, heteroaryl, heterosubstituted heteroaryl, RSG_1 or RSG_2 ;

40 Z_2 , Z_7 , Z_9 and Z_{10} are independently hydrocarbon, heterosubstituted hydrocarbon, heteroaryl, or heterosubstituted heteroaryl;

RSG_1 is an electron-affinic moiety;


RSG_2 is $-L-(RSG_1)_n$;

45 L is a linker comprising a chain of 1 to 30 atoms in the chain, the atoms being selected from the group consisting of C, O, N, S, Si, and P; and

n is an integer greater than or equal to 1.

55

13. A compound corresponding to the structure:



wherein

M comprises ammonium or is a metal;

5 R_1 is hydrogen or hydroxy;

R_2 is $-OT_2$, $-OC(O)Z_2$, $-OC(O)OZ_2$, RSG_1 or RSG_2 ;

R_4 is $-OT_4$, $-OC(O)Z_4$, $-OC(O)OZ_4$, RSG_1 or RSG_2 ;


R_7 is RSG_1 or RSG_2 ;

10 R_9 is hydrogen, keto, $-OT_9$, $-OC(O)Z_9$, $-OC(O)OZ_9$, RSG_1 or RSG_2 ;

R_{10} is hydrogen, keto, $-OT_{10}$, $-OC(O)Z_{10}$, $-OC(O)OZ_{10}$, RSG_1 or RSG_2 ;

R_7 , R_9 , and R_{10} independently have the alpha or beta stereochemical configuration;

15 R_{13} is hydroxy, protected hydroxy, keto, MO- or



R_{14} is hydrogen, hydroxy, protected hydroxy, RSG_1 or RSG_2 ;


20 T_2 , T_4 , T_9 , and T_{10} are independently hydrogen or hydroxy protecting group;

X_1 is $-OX_6$;

X_2 is hydrogen, hydrocarbon, heterosubstituted hydrocarbon, heteroaryl, or heterosubstituted heteroaryl;

- X_3 and X_4 are independently hydrogen,
 25 hydrocarbon, heterosubstituted hydrocarbon, heteroaryl,
 heterosubstituted heteroaryl or RSG_1 ;
 X_5 is $-X_{10}$, $-OX_{10}$, $-SX_{10}$, or $-NX_8X_{10}$;
 X_6 is hydrogen, hydrocarbon, heterosubstituted
 hydrocarbon, heteroaryl, heterosubstituted heteroaryl,
 30 hydroxy protecting group or a functional group which
 increases the water solubility of the taxane derivative;
 X_8 is hydrogen, hydrocarbon, heterosubstituted
 hydrocarbon, RSG_1 or RSG_2 ;
 X_{10} is hydrocarbon, heterosubstituted
 35 hydrocarbon, heteroaryl, heterosubstituted heteroaryl,
 RSG_1 or RSG_2 ;
 Z_2 , Z_4 , Z_9 and Z_{10} are independently hydrocarbon,
 heterosubstituted hydrocarbon, heteroaryl, or
 heterosubstituted heteroaryl;
 40 RSG_1 is an electron-affinic moiety;
 RSG_2 is $-L-(RSG_1)_n$;
 L is a linker comprising a chain of 1 to 30
 atoms in the chain, the atoms being selected from the
 group consisting of C, O, N, S, Si, and P; and
 45 n is an integer greater than or equal to 1.

14. A compound corresponding to the structure:



wherein

- 5 M comprises ammonium or is a metal;
 R_1 is hydrogen or hydroxy;
 R_2 is $-OT_2$, $-OCOZ_2$, $-OCOOZ_2$, RSG_1 or RSG_2 ;
 R_4 is $-OT_4$, $-OCOZ_4$, $-OCOOZ_4$, RSG_1 or RSG_2 ;


R_7 is hydrogen, keto, $-OT_7$, $-OCOZ_7$, $-OCOOZ_7$, RSG_1 or RSG_2 ;

10 R_9 is RSG_1 or RSG_2 ;

R_{10} is hydrogen, keto, $-OT_{10}$, $-OCOZ_{10}$, $-OCOOZ_{10}$, RSG_1 or RSG_2 ;

R_7 , R_9 , and R_{10} independently have the alpha or beta stereochemical configuration;

15 R_{13} is hydroxy, protected hydroxy, keto, MO- or



R_{14} is hydrogen, hydroxy, protected hydroxy, RSG_1 or RSG_2 ;

20 T_2 , T_4 , T_7 , and T_{10} are independently hydrogen or hydroxy protecting group;

X_1 is $-OX_6$;

X_2 is hydrogen, hydrocarbon, heterosubstituted hydrocarbon, heteroaryl, or heterosubstituted heteroaryl;

25 X_3 and X_4 are independently hydrogen, hydrocarbon, heterosubstituted hydrocarbon, heteroaryl, heterosubstituted heteroaryl or RSG_1 ;

X_5 is $-X_{10}$, $-OX_{10}$, $-SX_{10}$, or $-NX_8X_{10}$;

30 X_6 is hydrogen, hydrocarbon, heterosubstituted hydrocarbon, heteroaryl, heterosubstituted heteroaryl, hydroxy protecting group or a functional group which increases the water solubility of the taxane derivative;

X_8 is hydrogen, hydrocarbon, heterosubstituted hydrocarbon, RSG_1 or RSG_2 ;

35 X_{10} is hydrocarbon, heterosubstituted hydrocarbon, heteroaryl, heterosubstituted heteroaryl, RSG_1 or RSG_2 ;

58

Z_2 , Z_4 , Z_7 and Z_{10} are independently hydrocarbon, heterosubstituted hydrocarbon, heteroaryl, or heterosubstituted heteroaryl;


40 RSG_1 is an electron-affinic moiety;

RSG_2 is $-L-(RSG_1)_n$;

L is a linker comprising a chain of 1 to 30 atoms in the chain, the atoms being selected from the group consisting of C, O, N, S, Si, and P; and

45 n is an integer greater than or equal to 1.

15. A compound corresponding to the structure:



wherein

M comprises ammonium or is a metal;

5 R_1 is hydrogen or hydroxy;

R_2 is $-OT_2$, $-OCOZ_2$, $-OCOOZ_2$, RSG_1 or RSG_2 ;

R_4 is $-OT_4$, $-OCOZ_4$, $-OCOOZ_4$, RSG_1 or RSG_2 ;


R_7 is hydrogen, keto, $-OT_7$, $-OCOZ_7$, $-OCOOZ_7$, RSG_1 or RSG_2 ;

10 R_9 is hydrogen, keto, $-OT_9$, $-OCOZ_9$, $-OCOOZ_9$, RSG_1 or RSG_2 ;

R_{10} is RSG_1 or RSG_2 ;

R_7 , R_9 , and R_{10} independently have the alpha or beta stereochemical configuration;

15 R_{13} is hydroxy, protected hydroxy, keto, MO- or



R_{14} is hydrogen, hydroxy, protected hydroxy, RSG₁ or RSG₂;

20 T_2 , T_4 , T_7 and T_9 are independently hydrogen or hydroxy protecting group;

X_1 is $-OX_6$;

X_2 is hydrogen, hydrocarbon, heterosubstituted hydrocarbon, heteroaryl, or heterosubstituted heteroaryl;

25 X_3 and X_4 are independently hydrogen, hydrocarbon, heterosubstituted hydrocarbon, heteroaryl, heterosubstituted heteroaryl or RSG₁;

X_5 is $-X_{10}$, $-OX_{10}$, $-SX_{10}$, or $-NX_8X_{10}$;

30 X_6 is hydrogen, hydrocarbon, heterosubstituted hydrocarbon, heteroaryl, heterosubstituted heteroaryl, hydroxy protecting group or a functional group which increases the water solubility of the taxane derivative;

X_8 is hydrogen, hydrocarbon, heterosubstituted hydrocarbon, RSG₁ or RSG₂;

35 X_{10} is hydrocarbon, heterosubstituted hydrocarbon, heteroaryl, heterosubstituted heteroaryl, RSG₁ or RSG₂;

Z_2 , Z_4 , Z_7 and Z_9 are independently hydrocarbon, heterosubstituted hydrocarbon, heteroaryl, or heterosubstituted heteroaryl;

40 RSG₁ is an electron-affinic moiety;


RSG₂ is $-L-(RSG_1)_n$;

L is a linker comprising a chain of 1 to 30 atoms in the chain, the atoms being selected from the group consisting of C, O, N, S, Si, and P; and

45 n is an integer greater than or equal to 1.

60

16. A compound corresponding to the structure:



wherein

M comprises ammonium or is a metal;

5 R_1 is hydrogen or hydroxy;

R_2 is $-OT_2$, $-OCOZ_2$, $-OCOOZ_2$, RSG_1 or RSG_2 ;

R_4 is $-OT_4$, $-OCOZ_4$, $-OCOOZ_4$, RSG_1 or RSG_2 ;


R_7 is hydrogen, keto, $-OT_7$, $-OCOZ_7$, $-OCOOZ_7$, RSG_1
or RSG_2 ;

10 R_9 is hydrogen, keto, $-OT_9$, $-OCOZ_9$, $-OCOOZ_9$, RSG_1
or RSG_2 ;

R_{10} is hydrogen, keto, $-OT_{10}$, $-OCOZ_{10}$, $-OCOOZ_{10}$,
 RSG_1 or RSG_2 ;

15 R_7 , R_9 , and R_{10} independently have the alpha or
beta stereochemical configuration;

R_{13} is hydroxy, protected hydroxy, keto, MO- or



R_{14} is RSG_1 or RSG_2 ;

20 T_2 , T_4 , T_7 , T_9 and T_{10} are independently hydrogen
or hydroxy protecting group;

X_1 is $-OX_6$;

X_2 is hydrogen, hydrocarbon, heterosubstituted
hydrocarbon, heteroaryl, or heterosubstituted heteroaryl;

61

- X_3 and X_4 are independently hydrogen,
25 hydrocarbon, heterosubstituted hydrocarbon, heteroaryl,
heterosubstituted heteroaryl or RSG_1 ;
 X_5 is $-X_{10}$, $-OX_{10}$, $-SX_{10}$, or $-NX_8X_{10}$;
 X_6 is hydrogen, hydrocarbon, heterosubstituted
hydrocarbon, heteroaryl, heterosubstituted heteroaryl,
30 hydroxy protecting group or a functional group which
increases the water solubility of the taxane derivative;
 X_8 is hydrogen, hydrocarbon, heterosubstituted
hydrocarbon, RSG_1 or RSG_2 ;
 X_{10} is hydrocarbon, heterosubstituted
35 hydrocarbon, heteroaryl, heterosubstituted heteroaryl,
 RSG_1 or RSG_2 ;
 Z_2 , Z_4 , Z_7 , Z_9 and Z_{10} are independently
hydrocarbon, heterosubstituted hydrocarbon, heteroaryl or
heterosubstituted heteroaryl;
40 RSG_1 is an electron-affinic moiety;
 RSG_2 is $-L-(RSG_1)_n$;
 L is a linker comprising a chain of 1 to 30
atoms in the chain, the atoms being selected from the
group consisting of C, O, N, S, Si, and P; and
45 n is an integer greater than or equal to 1.

17. A method of killing tumor cells in a warm-blooded animal, the method comprising:

- (a) administering to the warm-blooded animal a
taxane containing an electron-affinic radiosensitizing
5 functional group,
(b) followed by, after a time interval
sufficient to enhance radiosensitization of the tumor
cells, irradiating the tumor cells with a dose of
radiation effective to kill the tumor cells.

18. A method as set forth in claim 16 further
comprising heat treating the tumor cells.

AMENDED CLAIMS


[received by the International Bureau on 4 February 1997((04.02.97); original claims 4 and 11-16 amended; remaining claims unchanged (7 pages)]

1. A compound comprising a taxane containing at least 2 electron-affinic radiosensitizing functional groups.

2. The compound of claim 1 wherein at least one of the radiosensitizing groups is a nitro-substituted carbocyclic or heterocyclic aromatic moiety which is attached to the C2, C4, C7, C9, C10 or C14 position of the taxane.

3. The compound of claim 1 wherein said radiosensitizing groups are independently selected from nitro-substituted carbocyclic and heterocyclic aromatic moieties and wherein at least one of said radiosensitizing groups is attached to the C2, C4, C7, C9, C10 or C14 position of the taxane.

4. A compound corresponding to the structure:



wherein

M comprises ammonium or is a metal;

5 R_1 is hydrogen or hydroxy;

R_2 is $-OT_2$, $-OCOT_2$, $-OCOT_2$, RSG_1 or RSG_2 ;

R_4 is $-OT_4$, $-OCOT_4$, $-OCOT_4$, RSG_1 or RSG_2 ;

R_7 is hydrogen, halogen, $-OT_7$, $-OCOT_7$, $-OCOT_7$, RSG_1 or RSG_2 ;

10 R_9 is hydrogen, keto, $-OT_9$, $-OCOT_9$, $-OCOT_9$, RSG_1 or RSG_2 ;

L is a linker comprising a chain of 1 to 10 atoms in the chain, the atoms being selected from the group consisting of C, O, N, S, Si, and P; and


35 n is 1 or 2.

8. The compound of claim 4 wherein RSG_1 is a heterocyclic aromatic moiety containing two or more heteroatoms.

9. The compound of claim 4 wherein RSG_1 is a metal complex.

10. The compound of claim 4 wherein RSG_1 is selected from the group consisting imidazoles, triazoles, pyridines, benzamides, furans, thiophenes, oxazoles and thiozoles possessing one or more nitro groups.

11. A compound corresponding to the structure:



wherein

M comprises ammonium or is a metal;

5 R_1 is hydrogen or hydroxy;

R_2 is RSG_1 or RSG_2 ;

R_4 is $-OT_4$, $-OCOZ_4$, $-OCOOZ_4$, RSG_1 or RSG_2 ;

R_7 is hydrogen, halogen, $-OT_7$, $-OCOZ_7$, $-OCOOZ_7$, RSG_1 or RSG_2 ;


10 R_9 is hydrogen, keto, $-OT_9$, $-OCOZ_9$, $-OCOOZ_9$, RSG_1 or RSG_2 ;

R_{10} is hydrogen, keto, $-OT_{10}$, $-OCOZ_{10}$, $-OCOOZ_{10}$, RSG_1 or RSG_2 ;

L is a linker comprising a chain of 1 to 30 atoms in
 45 the chain, the atoms being selected from the group
 consisting of C, O, N, S, Si, and P; and

n is an integer greater than or equal to 1.

12. A compound corresponding to the structure:



wherein

M comprises ammonium or is a metal;

5 R_1 is hydrogen or hydroxy;

R_2 is $-OT_2$, $-OC(O)Z_2$, $-OC(O)OZ_2$, RSG_1 or RSG_2 ;

R_4 is RSG_1 or RSG_2 ;


R_7 is hydrogen, halogen, $-OT_7$, $-OC(O)Z_7$, $-OC(O)OZ_7$, RSG_1
 or RSG_2 ;

10 R_9 is hydrogen, keto, $-OT_9$, $-OC(O)Z_9$, $-OC(O)OZ_9$, RSG_1 or
 RSG_2 ;

R_{10} is hydrogen, keto, $-OT_{10}$, $-OC(O)Z_{10}$, $-OC(O)OZ_{10}$, RSG_1 or
 RSG_2 ;


R_7 , R_9 , and R_{10} independently have the alpha or beta
 15 stereochemical configuration;

R_{13} is hydroxy, protected hydroxy, keto, MO- or




R_{14} is hydrogen, hydroxy, protected hydroxy, RSG_1 or
 RSG_2 ;

13. A compound corresponding to the structure:



wherein


- 5 M comprises ammonium or is a metal;
 R_1 is hydrogen or hydroxy;
 R_2 is $-OT_2$, $-OCOZ_2$, $-OCOOZ_2$, RSG_1 or RSG_2 ;
 R_4 is $-OT_4$, $-OCOZ_4$, $-OCOOZ_4$, RSG_1 or RSG_2 ;
 R_7 is RSG_1 or RSG_2 ;
 10 R_9 is hydrogen, keto, $-OT_9$, $-OCOZ_9$, $-OCOOZ_9$, RSG_1 or RSG_2 ;
 R_{10} is hydrogen, keto, $-OT_{10}$, $-OCOZ_{10}$, $-OCOOZ_{10}$, RSG_1 or RSG_2 ;
 R_7 , R_9 , and R_{10} independently have the alpha or beta stereochemical configuration;
 15 R_{13} is hydroxy, protected hydroxy, keto, MO- or



- R_{14} is hydrogen, hydroxy, protected hydroxy, RSG_1 or RSG_2 ;
 20 T_2 , T_4 , T_9 and T_{10} are independently hydrogen or hydroxy protecting group;
 X_1 is $-OX_6$;
 X_2 is hydrogen, hydrocarbon, heterosubstituted hydrocarbon, heteroaryl, or heterosubstituted heteroaryl;

- 25 X_3 and X_4 are independently hydrogen, hydrocarbon, heterosubstituted hydrocarbon, heteroaryl, heterosubstituted heteroaryl or RSG_1 ;
 X_5 is $-X_{10}$, $-OX_{10}$, $-SX_{10}$, or $-NX_8X_{10}$;
 X_6 is hydrogen, hydrocarbon, heterosubstituted
30 hydrocarbon, heteroaryl, heterosubstituted heteroaryl, hydroxy protecting group or a functional group which increases the water solubility of the taxane derivative;
 X_8 is hydrogen, hydrocarbon, heterosubstituted hydrocarbon, RSG_1 or RSG_2 ;
35 X_{10} is hydrocarbon, heterosubstituted hydrocarbon, heteroaryl, heterosubstituted heteroaryl, RSG_1 or RSG_2 ;
 Z_2 , Z_4 , Z_9 and Z_{10} are independently hydrocarbon, heterosubstituted hydrocarbon, heteroaryl, or heterosubstituted heteroaryl;
40 RSG_1 is an electron-affinic moiety;
 RSG_2 is $-L-(RSG_1)_n$;
 L is a linker comprising a chain of 1 to 30 atoms in the chain, the atoms being selected from the group consisting of C, O, N, S, Si, and P; and
45 n is an integer greater than or equal to 1.

14. A compound corresponding to the structure:



wherein

- M comprises ammonium or is a metal;
5 R_1 is hydrogen or hydroxy;
 R_2 is $-OT_2$, $-OC(O)Z_2$, $-OC(O)OZ_2$, RSG_1 or RSG_2 ;
 R_4 is $-OT_4$, $-OC(O)Z_4$, $-OC(O)OZ_4$, RSG_1 or RSG_2 ;

Z_2 , Z_4 , Z_7 and Z_{10} are independently hydrocarbon, heterosubstituted hydrocarbon, heteroaryl, or
 40 heterosubstituted heteroaryl;


RSG_1 is an electron-affinic moiety;

RSG_2 is $-L-(RSG_1)_n$;

L is a linker comprising a chain of 1 to 30 atoms in the chain, the atoms being selected from the
 45 group consisting of C, O, N, S, Si, and P; and

n is an integer greater than or equal to 1.

15. A compound corresponding to the structure:



wherein

M comprises ammonium or is a metal;

5 R_1 is hydrogen or hydroxy;

R_2 is $-OT_2$, $-OCOZ_2$, $-OCOOZ_2$, RSG_1 or RSG_2 ;

R_4 is $-OT_4$, $-OCOZ_4$, $-OCOOZ_4$, RSG_1 or RSG_2 ;


R_7 is hydrogen, keto, $-OT_7$, $-OCOZ_7$, $-OCOOZ_7$, RSG_1
 or RSG_2 ;

10 R_9 is hydrogen, keto, $-OT_9$, $-OCOZ_9$, $-OCOOZ_9$, RSG_1
 or RSG_2 ;


R_{10} is RSG_1 or RSG_2 ;

R_7 , R_9 , and R_{10} independently have the alpha or beta stereochemical configuration;

15 R_{13} is hydroxy, protected hydroxy, keto, MO- or



16. A compound corresponding to the structure:



wherein

M comprises ammonium or is a metal;

5

R_1 is hydrogen or hydroxy;

R_2 is $-OT_2$, $-OCOT_2$, $-OCOT_2$, RSG_1 or RSG_2 ;

R_4 is $-OT_4$, $-OCOT_4$, $-OCOT_4$, RSG_1 or RSG_2 ;

R_7 is hydrogen, keto, $-OT_7$, $-OCOT_7$, $-OCOT_7$, RSG_1

or RSG_2 ;

10

R_9 is hydrogen, keto, $-OT_9$, $-OCOT_9$, $-OCOT_9$, RSG_1


or RSG_2 ;

R_{10} is hydrogen, keto, $-OT_{10}$, $-OCOT_{10}$, $-OCOT_{10}$,

RSG_1 or RSG_2 ;

15 R_7 , R_9 , and R_{10} independently have the alpha or
beta stereochemical configuration;

R_{13} is hydroxy, protected hydroxy, keto, MO- or



;

R_{14} is RSG_1 or RSG_2 ;


20 T_2 , T_4 , T_7 , T_9 and T_{10} are independently hydrogen
or hydroxy protecting group;

X_1 is $-OX_6$;

X_2 is hydrogen, hydrocarbon, heterosubstituted
hydrocarbon, heteroaryl, or heterosubstituted heteroaryl;


1/18

FIG. 1




2/18

FIG. 2




3/18

FIG. 3




4/18

FIG. 4




5/18

FIG. 5




6/18

FIG. 6




7/18

FIG. 7




8/18

FIG. 8




9/18

FIG. 9




10/18

FIG. 10




11/18

FIG. 11




12/18

FIG.12




13/18

FIG. 13




14/18

FIG. 14




15/18

FIG. 15




16/18

FIG. 16




17/18

FIG. 17



18/18

FIG. 18



INTERNATIONAL SEARCH REPORT

International application No.
PCT/US96/14613

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : A61K 31/335; C07D 305/14

US CL : 514/449; 549/510, 511

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/449; 549/510, 511

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
NONEElectronic data base consulted during the international search (name of data base and, where practicable, search terms used)
none

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X, P	US 5,466,834 A (HOLTON) 14 November 1995, see the entire document.	1-18



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Z* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

22 NOVEMBER 1996

Date of mailing of the international search report

05 DEC 1996

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

BA K. TRINH

Telephone No. (703) 308-1235